

MMA2024

27th International Conference
**Mathematical Modelling
and Analysis**

May 28 - 31, 2024
Pärnu, Estonia

**CONFERENCE PROGRAMME &
ABSTRACTS**

Conference Organizers

Institute of Mathematics and Statistics of University of Tartu (Estonia)

Vilnius Gediminas Technical University (Lithuania)

Estonian Mathematical Society

International Program Committee

S. Asmuss (Latvia), L. Brugnano (Italy), I. Bula (Latvia), R. Čiegis (Lithuania), N. J. Ford (UK), F. Gaspar (Spain), A. Gritsans (Latvia), U. Hämarik (Estonia), O. Iliev (Germany), Z. Jackiewicz (USA), J. Janno (Estonia), A. Kolishkins (Latvia), A. Pedas (Estonia), K. Pileckas (Lithuania), M. Radziunas (Germany), M. Ragulskis (Lithuania), A. Reinfelds (Latvia), F. Sadyrbaev (Latvia), M. Sapagovas (Lithuania), A. Sostaks (Latvia), V. Starikovicius (Lithuania), A. Stikonas (Lithuania), U. Strautins (Latvia), G. Vainikko (Estonia)

Organizing Committee

A. Pedas (Chairman), R. Čiegis (Vice-chairman), T. Raus (Vice-chairman), U. Hämarik, U. Kangro, E. Kiriaed, M. Lillemäe, K. Lätt, K. Orav-Puurand, H. B. Soots, M. Vikerpuur

Supported by

Estonian Research Council Grant PRG864

Estonian Mathematical Society

PLENARY SPEAKERS

Luigi Brugnano (Università di Firenze, Italy)

Recent advances in the numerical solution of ordinary and fractional differential equations

Raimondas Čiegis (Vilnius Gediminas technical university, Lithuania)

LONG STANDING and NEVER OLD QUESTION: WHO is a BOSS (STABILITY or APPROXIMATION)

Kai Diethelm (Technical University of Applied Sciences Würzburg-Schweinfurt, Germany)

Fast Numerical Algorithms for the Evaluation of Riemann-Liouville Integrals

Neville J Ford (University of Chester, United Kingdom)

Modelling and simulation of systems with complex characteristics: certainties, uncertainties and compromises

Zdzisław Jackiewicz (Arizona State University, United States)

Local error estimation for implicit-explicit general linear methods

Pedro Lima (University of Lisbon, Portugal)

Mathematical Modeling of Neural Fields with Diffusion

Minvydas Ragulskis (Kaunas University of Technology, Lithuania)

Solitary solutions to the Hepatitis C model with the proliferation of infected and uninfected hepatocytes

Tarmo Soomere (Tallinn University of Technology, Estonia)

Mathematics of extremes in coastal and marine science

Martin Stynes (Beijing Computational Science Research Center, China)

A general collocation analysis for weakly singular Volterra integral equations with variable exponent

MINI-SYMPOSIUMS

Application of zeta functions for approximation of analytical functions

Organized by **A. Laurinčikas** and **D. Šiaučiūnas**

Fractional calculus, fractal sets and zeta functions

Organized by **E. Guariglia**

*Nonlocal differential problems and their applications**

Organized by **K. Kaulakytė** and **R. Čiegis**

*This mini-symposium is supported by the Research Council of Lithuania (LMTLT), agreement No. S-MIP-23-43.

LIST OF TALKS

LOCAL FRACTAL FUNCTIONS OF HIGHER ORDER AND SYMMETRIES	9
Waldo Arriagada	9
ON INTERACTION OF SETS OF PERIODIC SOLUTIONS	10
Svetlana Atslega, Felix Sadyrbaev	10
EXPLICIT TIME INTEGRATION OF SOURCE-FREE DYNAMICAL SYSTEMS	11
Jānis Bajārs, Anna Simina	11
ON THE LAPLACE TRANSFORM OF THE RIEMANN ZETA-FUNCTION II	12
Aidas Balčiūnas	12
INVESTIGATION OF A DISCRETE STURM–LIOUVILLE PROBLEM WITH TWO-POINT NONLOCAL BOUNDARY CONDITION	13
Kristina Bingelė, Artūras Štikonas	13
HIGH ORDER SECOND DERIVATIVE DIAGONALLY IMPLICIT MULTISTAGE INTEGRATION METHODS FOR ODES	14
Michał Braś	14
RECENT ADVANCES IN THE NUMERICAL SOLUTION OF ORDINARY AND FRACTIONAL DIFFERENTIAL EQUATIONS	15
Luigi Brugnano	15
HYERS-ULAM STABILITY OF IMPLICIT VOLTERRA EQUATIONS ON TIME SCALES	16
Andrejs Reinfelds, Shraddha Ramanbhai Christian	16
LONG STANDING AND NEVER OLD QUESTION: WHO IS A BOSS (STABILITY OR APPROXIMATION)	17
Raimondas Čiegis	17
MODELLING OF BACTERIAL PLUME FORMATION IN A CIRCULAR CON- TAINER	18
Boleslovas Dapkūnas	18
FAST NUMERICAL ALGORITHMS FOR THE EVALUATION OF RIEMANN- LIOUVILLE INTEGRALS	19
Kai Diethelm	19
CO AND CO₂ CONCENTRATION IN BIOMASS COMBUSTION PROCESS DE- PENDENT ON BIOMASS POROSITY	20
Maris Gunars Dzenis	20
A GLOBAL APPROXIMATION METHOD FOR SECOND-KIND VOLTERRA- FREDHOLM EQUATIONS	21
Luisa Fermo, Domenico Mezzanotte, Donatella Occorsio	21
INTEGRATING FRACTIONAL DIFFERENTIAL EQUATIONS WITH NEURAL NETWORKS FOR DYNAMICAL SYSTEMS MODELLING	22
Cecília Coelho, M. Fernanda P. Costa, Luís L. Ferrás	22
MODELLING AND SIMULATION OF SYSTEMS WITH COMPLEX CHARAC- TERISTICS: CERTAINTIES, UNCERTAINTIES AND COMPROMISES	23

Neville J Ford	23
A NONLINEAR BACKWARD DIFFUSION ALGORITHM FOR IMAGE RESTORATION	24
Lorella Fatone, Daniele Funaro	24
ON DISCRETE SHIFTS OF THE MELLIN TRANSFORM OF THE RIEMMAN ZETA-FUNCTION	25
Virginija Garbaliuskienė, Antanas Laurinčikas, Darius Šiaučiūnas	25
NUMERICAL APPROACHES TO SOLVE AN INVERSE PROBLEM FOR A FRACTIONAL DIFFUSION-WAVE EQUATION WITH MULTI-TERM ...	26
Hany Gerges, Jaan Janno	26
JOINT MIXED LIMIT THEOREM FOR EPSTEIN AND HURWITZ ZETA-FUNCTIONS	27
Hany Gerges, Renata Macaitienė	27
FRACTIONAL CALCULUS OF ZETA FUNCTIONS	28
Emanuel Guariglia	28
LOCAL ERROR ESTIMATION FOR IMPLICIT-EXPLICIT GENERAL LINEAR METHODS	29
Zdzisław Jackiewicz	29
APPROXIMATION OF ANALYTIC FUNCTIONS BY SHIFTS OF ABSOLUTELY CONVERGENT DIRICHLET SERIES RELATED TO PERIODIC ...	30
Mindaugas Jasas	30
REGULARITY RESULTS FOR VERY WEAK TIME-PERIODIC POISEUILLE-TYPE SOLUTION WITH MINIMALLY REGULAR FLOW RATE	31
Rita Juodagalvytė	31
NOTES ON SIMULTANEOUS APPROXIMATION BY THE CLASS OF ZETA-FUNCTIONS	32
Roma Kačinskaitė	32
STALLING IN QUEUING SYSTEMS WITH HETEROGENEOUS CHANNELS	33
Liudvikas Kaklauskas	33
A NONLINEAR MATHEMATICAL MODEL OF THE COURSE AND THERAPY OF RHEUMATOID ARTHRITIS AND ITS REALIZATION	34
Zviad Kalichava, Kakhaber Odisharia, Vladimer Odisharia, Nona Janikashvili	34
LEARNING HAMILTONIAN DYNAMICS WITH STRUCTURE-PRESERVING NEURAL NETWORKS AND DIMENSIONALITY REDUCTION	35
Dāvis Kalvāns, Jānis Bajārs	35
ON CONVERGENCE CONDITIONS IN SELF-REGULARIZATION OF ILL-POSED PROBLEMS BY PROJECTION METHODS	36
Urve Kangro, Uno Hämarik	36
TIME PERIODIC INVERSE PROBLEM WITH NONLOCAL CONDITION	37
Kristina Kaulakytė	37
PERIODIC CUBIC SPLINE HISTOPOLATION	38

Evely Kirsiaed, Peeter Oja	38
POINTWISE-IN-TIME A-PRIORI AND A-POSTERIORI ERROR CONTROL FOR TIME-FRACTIONAL PARABOLIC EQUATIONS	39
Natalia Kopteva	39
CURVE ESTIMATION WITH MODIFIED COMPLETE SPLINES AND EXPO- NENTIAL PARAMETERIZATION	40
Ryszard Kozera, Magdalena WilkoŁazka	40
BLOOD FLOW MODELLING IN HUMAN HEART USING FSI	41
Nikolajus Kozulinas	41
ABOUT THE NUMERICAL SOLUTION OF SINGULAR FRACTIONAL INTEGRO-DIFFERENTIAL EQUATIONS	42
Kaido Lätt	42
APPROXIMATION BY BEURLING'S ZETA-FUNCTIONS	43
Antanas Laurinćikas	43
ON THE NUMERICAL SOLUTION OF THE NEUMANN PROBLEM FOR LAPLACE'S EQUATION IN PLANAR DOMAINS WITH CORNERS	44
Peter Junghanns, Concetta Laurita	44
ON CENTRAL PART INTERPOLATION APPROXIMATIONS FOR SYSTEMS OF FRACTIONAL DIFFERENTIAL EQUATIONS	45
Margus Lillemäe	45
MATHEMATICAL MODELING OF NEURAL FIELDS WITH DIFFUSION	46
Daniele Avitabile, Nikolai Chemetov, Pedro Lima	46
SOME GENERALIZED LIMIT THEOREMS FOR THE EPSTEIN ZETA-FUNCTION	47
Renata Macaitienė	47
ON MATHEMATICAL MODELLING AND SIMULATION OF IMMUNE SYS- TEM'S RESPONSE IN TUMOR TREATMENT	48
Maksims Marinaki, Manfreds Šnepš-Šneppe	48
ON GENERALIZED SHIFTS OF THE LERCH ZETA-FUNCTION	49
Toma Mikalauskaitė	49
STABILITY ANALYSIS OF STATIONARY MOTION OF THE 3D SWINGING ATWOOD'S MACHINE	50
Alexander Prokopenya	50
ON THE CONVERGENCE OF THE DIFFERENCE SCHEME FOR NONLIN- EAR PARABOLIC EQUATION WITH INTEGRAL BOUNDARY ...	51
K. Pupalaigė, M. Sapagovas	51
SOLITARY SOLUTIONS TO THE HEPATITIS C MODEL WITH THE PROLIF- ERATION OF INFECTED AND UNINFECTED HEPATOCYTES	52
Minvydas Ragulskis	52
NEW HEURISTIC PARAMETER CHOICE FOR THE CLASS OF REGULAR- IZATION METHODS	53

Toomas Raus	53
A POLYNOMIAL COLLOCATION METHOD FOR THE LIDTHILL'S PROBLEM	54
Magda Rebelo	54
DISCRETE VERSION OF THE MISHOU THEOREM FOR ABSOLUTELY CONVERGENT DIRICHLET SERIES RELATED TO PERIODIC ZETA-...	55
Aidas Balčiūnas, Audrone Rimkevičienė	55
A GLOBAL APPROXIMATION METHOD FOR SECOND-KIND NONLINEAR INTEGRAL EQUATIONS	56
Luisa Fermo, Anna Lucia Laguardia, Concetta Laurita, Maria Grazia Russo	56
ON SENSITIVE DEPENDENCE OF SOLUTIONS ON THE INITIAL DATA IN ORDINARY DIFFERENTIAL EQUATIONS	57
Felix Sadyrbaev	57
ON THE MEAN SQUARE OF THE HURWITZ ZETA-FUNCTION IN SHORT INTERVALS	58
Darius Šiaučiūnas	58
ON ZETA-FUNCTION OF CERTAIN CUSP FORMS	59
Antanas Laurinčikas, Darius Šiaučiūnas	59
MATHEMATICS OF EXTREMES IN COASTAL AND MARINE SCIENCE	60
Tarmo Soomere	60
COLLOCATION BASED APPROXIMATIONS FOR FRACTIONAL BOUNDARY VALUE PROBLEMS	61
Hanna Britt Soots	61
DISCRETE STURM-LIOUVILLE PROBLEM FOR TWO-DIMENSIONAL ELLIPTIC EQUATION WITH THE MULTIPLE INTEGRAL IN NONLOCAL ...	62
Artūras Štikonas, Mifodijus Sapagovas	62
FINITE DIFFERENCE METHOD FOR ELLIPTIC EQUATION WITH TWO MULTIPLE INTEGRAL CONDITIONS	63
Olga Štikonienė, Mifodijus Sapagovas	63
ON SIMULATION WITH PARTICLE MOTION TRACKING FOR BIOMASS EXTRACTION PROCESS	64
Uldis Strautiņš, Maksims Marinaki	64
A GENERAL COLLOCATION ANALYSIS FOR WEAKLY SINGULAR VOLTERRA INTEGRAL EQUATIONS WITH VARIABLE EXPONENT	65
Martin Stynes	65
ON DERIVATIVE SAMPLING AND CONVERGENCE IN VARIATION OF GENERALIZED SHANNON SAMPLING SERIES	66
Gert Tamberg, Maria Zeltser	66
APPROXIMATION OF ANALYTIC FUNCTIONS BY SHIFTS OF THE PERIODIC ZETA-FUNCTIONS	67
Monika Tekorė	67

TIME-FREQUENCY ANALYSIS IN EUCLIDEAN SPACES	68
Ville Turunen	68
THE FUČÍK TYPE PROBLEM WITH NONLOCAL TWO-POINT BOUNDARY CONDITIONS	69
Sigita Urbonienė, Natalija Sergejeva	69
OPERATORS OF FUZZY MATHEMATICAL MORPHOLOGY	70
Ksenija Varfolomejeva, Alexander Šostak	70
APPROXIMATE SOLUTIONS TO LINEAR FRACTIONAL INTEGRO-DIFFERENTIAL EQUATIONS	71
Mikk Vikerpuur	71
ON A SOLVABILITY OF A THREE-PARAMETER PROBLEM ARISING IN HEAT TRANSFER	72
Armands Gritsans, Inara Yermachenko	72

LOCAL FRACTAL FUNCTIONS OF HIGHER ORDER AND SYMMETRIES

WALDO ARRIAGADA

Wenzhou-Kean University

88 Daxue, Ouhai, Wenzhou, Zhejiang Province, China, 325060.

E-mail: warriaga@kean.edu

In this talk we prove the existence of local fractal functions of the Orlicz-Sobolev class of order $m \geq 0$. The graph of a local fractal function coincides with the attractor of an appropriate iterated function system (ifs), whose construction is fairly standard. Local fractal functions appear naturally as the fixed points of the Read-Bajraktarević operator when restricted to a suitable Orlicz-Sobolev space. Our results extend some of the outcomes obtained by Massopust on Lebesgue and Sobolev spaces to higher order, dimension and function spaces (where the role of the norm is now played by a Young function). As an application, we discuss the existence of local fractal functions on analytic spaces and we highlight the existence of symmetries arising from the complex conjugation.

REFERENCES

- [1] Arriagada W., *Local fractal functions on Orlicz-Sobolev spaces*, preprint (2022).
- [2] Arriagada W., *Matuszewska-Orlicz indices of the Sobolev conjugate Young function*. Partial Differential Equations in Applied Mathematics, vol. **3** (2021).
- [3] Bajraktarević M., *Sur une équation fonctionnelle*, Glasnik Mat. Fiz. I Astr. **12**, pp 201–205, 1957.
- [4] Barnsley M., Hegland M., and Massopust P., *Numerics and fractals*, Bulletin of the Institute of Mathematics Academia Sinica (New Series), vol. **9**, No. 3, 389–430, 2014.
- [5] Constantine G.M., and Savits T.H., *A multivariate Faà di Bruno formula with applications*, Transactions of the American Mathematical Society, **348**(2), 503–520, 1996.
- [6] Fukagai N., Ito M., and Narukawa K., *Positive solutions of Quasilinear Elliptic Equations with Critical Orlicz-Sobolev Nonlinearity on \mathbb{R}^N* , Funkcialaj Ekvacioj **49**, 235–267, 2006.
- [7] Gossez J. *Nonlinear elliptic boundary value problems for equations with rapidly (or slowly) increasing coefficients*, Trans. Amer. Math. Soc., **190**, 163–205, 1974.
- [8] Huentutripay J., and Manásevich R. *Nonlinear eigenvalues for a Quasilinear Elliptic System in Orlicz-Sobolev Spaces*, Journal of Dynamics and Differential Equations **18**, 901–921, 2006.
- [9] Maligranda L., *Indices and interpolation*. Seminars in Math., Univ. of Campinas, Campinas SP: Brazil; 81–95, 1989.
- [10] Massopust P.R., *Fractal Functions, Fractal Surfaces, and Wavelets*, Academic Press, Inc., San Diego, CA, 1994.
- [11] McClure M., *The Read-Bajraktarević Operator*, Mathematica in Education and Research, Vol. 11, No. 3, 356–362, (2006).
- [12] Vrscay E.R., *Moment and Collage Methods for the Inverse Problem of Fractal Construction with Iterated Function Systems*. In: Peitgen HO, Henriques JM, Penedo LF, eds. Fractals in the Fundamental and Applied Sciences. North-Holland, 443–461. 1991.

ON INTERACTION OF SETS OF PERIODIC SOLUTIONS

SVETLANA ATSLEGA^{1,2}, FELIX SADYRBAEV^{1,3}

Institute of Mathematics and Computer Science, University of Latvia

Rainis boulevard 29, LV-1459 Rīga, Latvia

Latvia University of Life Sciences and Technologies, Institute of Mathematics and Physics

2 Lielā street, Jelgava, LV-3001, Latvia

Daugavpils University

13 Vienības street, Daugavpils, LV-5401, Latvia

E-mail: svetlana.atslega@lbtu.lv

A class of autonomous ordinary differential equations of the second order is considered, which is characterized by the existence of several period annuli (the sets of periodic solutions).

The number of solutions to the Neumann boundary value problem is studied. The bifurcation diagrams are constructed. The estimates of the number of solutions are obtained.

Perturbations of the autonomous equations with period annuli by external periodic force are studied. The chaotic behavior of solutions to the perturbed equation is observed. Visual evidences of chaotic behavior are presented. The Lyapunov exponents are used for the analysis of the sensitive dependence of solutions on the initial data.

REFERENCES

- [1] S. Atslega and F. Sadyrbaev. On periodic solutions of Liénard type equations. *Mathematical Modelling and Analysis*, **18** (5):708–716, 2013.
- [2] Y. Kozmina and F. Sadyrbaev. On a Maximal Number of Period Annuli. *Abstract and Applied Analysis*, **2011** Article ID 393875, 8 pages.
- [3] J.C. Sprott. *Elegant Chaos: Algebraically Simple Chaotic Flows*. World Scientific, 2010.

EXPLICIT TIME INTEGRATION OF SOURCE-FREE DYNAMICAL SYSTEMS

JĀNIS BAJĀRS, ANNA SIMINA

Faculty of Physics, Mathematics and Optometry, University of Latvia

Jelgavas street 3, Riga LV-1004, Latvia

E-mail: janis.bajars@lu.lv, as16069@students.lu.lv

The flow ϕ_t of a source-free dynamical system $\dot{x} = f(x)$, i.e., with a divergence-free vector field: $\operatorname{div} f(x) = 0$, is phase volume-preserving: $\det \frac{\partial \phi_t(x)}{\partial x} = 1$ for all t and x , a desirable property to be maintained by a numerical time integration method. Feng and Shang have shown in [1] that every divergence-free vector field $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$ can be written as the sum of $n - 1$ vector fields $f = f_{1,2} + f_{2,3} + \dots + f_{n-1,n}$, where each of $f_{k,k+1}$ is Hamiltonian in the variables (x_k, x_{k+1}) , i.e., there exist functions $H_{k,k+1} : \mathbb{R}^n \rightarrow \mathbb{R}$ such that $f_{k,k+1} = \left(0, \dots, 0, -\frac{\partial H_{k,k+1}}{\partial x_{k+1}}, \frac{\partial H_{k,k+1}}{\partial x_k}, 0, \dots, 0\right)^T$. Considering this decomposition, a phase volume-preserving numerical method is obtained by applying a splitting method with symplectic substeps, which are volume-preserving. Different splitting approaches allow the construction of symmetric as well as higher-order volume-preserving methods [2]. In general, the obtained methods are one-dimensionally implicit unless $\frac{\partial f_i(x)}{\partial x_i} = 0$ for all $i = 1, \dots, n$.

To address the question of fully explicit symplectic integration of general (nonseparable) Hamiltonian dynamics, Tao in [3] proposed symplecticity-preserving time integration methods in an extended phase space by introducing a mechanical restraint that binds together two copies of the original phase space. In [3], good long-time numerical simulation properties were demonstrated, and an error bound for integrable dynamics was derived. In this work, we propose explicit time integration methods of source-free dynamical systems in an extended phase space, similarly to [3], by considering the augmented Hamiltonians: $\bar{H}_{k,k+1}(x_1, \dots, x_n, y_1, \dots, y_n) = H_{k,k+1}^A(x_1, \dots, x_{k-1}, y_k, x_{k+1}, \dots, x_n) + H_{k,k+1}^B(x_1, \dots, x_k, y_{k+1}, x_{k+2}, \dots, x_n) + \frac{1}{2}\omega((x_k - y_k)^2 + (x_{k+1} - y_{k+1})^2)$, where ω is a binding constant of system's variable x with its copy y . With augmented Hamiltonians, we obtain vector fields

$$\bar{f}_{k,k+1} = \left(0, \dots, 0, -\frac{\partial H_{k,k+1}^A}{\partial x_{k+1}} - \omega(x_{k+1} - y_{k+1}), \frac{\partial H_{k,k+1}^B}{\partial x_k} + \omega(x_k - y_k), 0, \dots, 0, \right. \\ \left. 0, \dots, 0, -\frac{\partial H_{k,k+1}^B}{\partial y_{k+1}} + \omega(x_{k+1} - y_{k+1}), \frac{\partial H_{k,k+1}^A}{\partial y_k} - \omega(x_k - y_k), 0, \dots, 0\right)^T,$$

which allow the construction of explicit splitting methods for general source-free dynamical systems. The properties of proposed explicit numerical methods are explored and numerically demonstrated.

REFERENCES

- [1] K. Feng and Z. Shang. Volume-preserving algorithms for source-free dynamical systems. *Numerische Mathematik*, **71** 451–463, 1995.
- [2] E. Hairer and C. Lubich, and G. Wanner. *Geometric Numerical Integration: Structure-preserving Algorithms for Ordinary Differential Equations*. Springer Science & Business Media, 2006.
- [3] M. Tao. Explicit symplectic approximation of nonseparable Hamiltonians: Algorithm and long time performance. *Physical Review E*, **94** 043303, 2016.

ON THE LAPLACE TRANSFORM OF THE RIEMANN ZETA-FUNCTION II

AIDAS BALČIŪNAS

Institute of Mathematics, Faculty of Mathematics and Informatics, Vilnius University

Naugarduko str. 24, LT-03225 Vilnius, Lithuania

E-mail: `aidas.balciunas@mif.vu.lt`

The present talk is devoted to the Laplace transform

$$\mathfrak{L}(s, |\zeta|^{2k}) = \int_0^\infty \left| \zeta \left(\frac{1}{2} + ix \right) \right|^{2k} e^{-sx} dx$$

of the Riemann zeta-function $\zeta(s)$ with complex variable s , where $k \in \mathbb{N}$. In [1], only the case $k = 2$ has been considered. In [2] we had the case with the odd k . This report is a more general case of the transform for arbitrary k using the \mathfrak{G} Meijer function.

REFERENCES

- [1] A.Ivič. The Laplace transform of the fourth moment of the zeta-function. *Univ. Beograd. Publ. Elektrotehn. Scr. Mat.*, **11** 41–48, 2000.
- [2] A. Balčiūnas. On the Laplace transform of the Riemann zeta-function. In: *26 th International Conference on Mathematical Modeling and Analysis (MMA 2023)*, 6, 2023.

INVESTIGATION OF A DISCRETE STURM–LIOUVILLE PROBLEM WITH TWO-POINT NONLOCAL BOUNDARY CONDITION

KRISTINA BINGELĖ¹ and ARTŪRAS ŠTIKONAS²

¹*Vilnius Gediminas Technical University*

Saulėtekio al. 11, LT-10223 Vilnius, Lithuania

²*Institute of Applied Mathematics, Vilnius University*

Naugarduko str. 24, LT-03225 Vilnius, Lithuania

E-mail: kristina.bingele@vilniustech.lt; arturas.stikonas@mif.vu.lt

We investigate SLP with one classical Dirichlet Boundary Condition (BC) or Neumann BC:

$$\begin{aligned} -u'' &= \lambda u, & t \in (0, 1), & \lambda \in \mathbb{C}, & (1) \\ u(0) &= 0 & \text{or} & & u'(0) = 0, & (2d, n) \end{aligned}$$

and another two-points Nonlocal Boundary Condition (NBC)

$$u(1) = \gamma u(\xi), \quad u'(1) = \gamma u'(\xi), \quad u(1) = \gamma u'(\xi) \quad \text{or} \quad u'(1) = \gamma u(\xi), \quad (3_{1,2,3,4})$$

where NBC's parameter $\gamma \in \mathbb{R}$ and $\xi \in [0, 1]$.

We introduce a uniform grids in $[0, 1]$: $\bar{\omega}^h = \{t_j = jh, j = \overline{0, n}\}$, $\omega^h = \{t_j = jh, j = \overline{1, n-1}\}$ with stepsizes $h_j \equiv h$ and $\omega_{1/2}^h = \{t_{j+1/2} = (t_j + t_{j+1})/2, j = \overline{0, n-1}\}$ with stepsizes $h_{j+1/2} = t_{j+1/2} - t_{j-1/2} \equiv h$. Additionally, we use a nonuniform grid $\bar{\omega}_{1/2}^h = \omega_{1/2}^h \cup \{t_{-1/2} = 0, t_{n+1/2} = n\}$ where stepsizes $h_{1/2} = t_{1/2} - t_{-1/2} = h/2$, $h_{n+1/2} = t_{n+1/2} - t_{n-1/2} = h/2$. We make an assumption that $\xi = m/n$ is located on the grid $\bar{\omega}^h$. We approximate differential SLP (1)–(3_{1,2,3,4}) by the discrete SLP, using natural approximation of derivative $\bar{\delta}$:

$$\begin{aligned} -\delta^2 U &= \lambda U, & t \in \omega^h, & \lambda \in \mathbb{C}_\lambda, & (4) \\ U_0 &= 0 & \text{or} & & (\bar{\delta}U)_0 = 0, & (5d, n) \\ U_n &= \gamma U_m, & (\bar{\delta}U)_n &= \gamma(\bar{\delta}U)_m, & U_n &= \gamma(\bar{\delta}U)_m \quad \text{or} \quad (\bar{\delta}U)_n = \gamma U_m, & (6_{1,2,3,4}) \end{aligned}$$

We investigate discrete SLP and analyze how complex eigenvalues of this problem depend on the parameters of the two-points NBC. Some results for the both SLP were presented in [1; 3].

REFERENCES

- [1] K. Bingelė, A. Bankauskienė and A. Štikonas. Investigation of spectrum for a Sturm–Liouville problem with with two-point nonlocal boundary conditions. *Math. Model. Anal.*, **25** (1):53–70, 2020.
- [2] A.A. Samarskii and E.S. Nikolaev. *Numerical Methods for Grid Equations*. Birkhäuser Verlag, Basel, Boston, Berlin, 1989. (Vol. I, *Iterative Methods*; Vol. II, *Direct Methods*)
- [3] K. Bingelė and A. Štikonas. Investigation of a Discrete Sturm–Liouville problem with Two-Point Nonlocal Boundary Condition and Natural Approximation of a Derivative in Boundary Condition. *Math. Model. Anal.*, **29** (2):309–330, 2024.

HIGH ORDER SECOND DERIVATIVE DIAGONALLY IMPLICIT MULTISTAGE INTEGRATION METHODS FOR ODES

MICHAŁ BRAŚ

AGH University

al. Mickiewicza 30, 30-059 Kraków, Poland

E-mail: bras@agh.edu.pl

For the numerical solution of the m -dimensional system of ordinary differential equations (ODEs)

$$y'(x) = f(y(x)), \quad y(x_0) = y_0, \quad x \in [x_0, \bar{x}],$$

we consider the class of second derivative diagonally implicit multistage integration methods (SDIM-SIMs). These methods are a specific subclass of second derivative general linear methods, which are represented in the following form:

$$Y_i^{[n]} = h \sum_{j=1}^s a_{ij} f(Y_j^{[n]}) + h^2 \sum_{j=1}^s \bar{a}_{ij} g(Y_j^{[n]}) + \sum_{j=1}^r u_{ij} y_j^{[n-1]}, \quad i = 1, 2, \dots, s,$$
$$y_i^{[n]} = h \sum_{j=1}^s b_{ij} f(Y_j^{[n]}) + h^2 \sum_{j=1}^s \bar{b}_{ij} g(Y_j^{[n]}) + \sum_{j=1}^r v_{ij} y_j^{[n-1]}, \quad i = 1, 2, \dots, r,$$

where $g(\cdot) = f'(\cdot)f(\cdot)$.

In [1] methods of order $p \leq 4$ were investigated. In this talk, we aim to extend this research by investigating methods of order $p \geq 5$ as presented in the [2]. Developing methods of higher order requires establishment of conditions based on the parameters of the methods. These conditions, in the form of a system of polynomial equations, cannot be obtained and solved through symbolic manipulation tools. Therefore, we propose an approach for constructing implicit and explicit SDIMSIM with Runge–Kutta stability property using a Fourier series method variation, previously utilized for constructing high-order general linear methods. Examples of fifth and sixth order explicit and implicit SDIMSIMs, suitable for both non-stiff and stiff differential systems in a sequential computing environment, are provided. Additionally, the effectiveness of the newly derived methods is validated through numerical experiments.

This is a joint work with Ali Abdi, Mohammad Sharifi and Gholamreza Hojjati from University of Tabriz, Iran.

REFERENCES

- [1] A. Abdi, M. Braś, and G. Hojjati. On the construction of second derivative diagonally implicit multistage integration methods for ODEs. *Applied Numerical Mathematics*, **76**, 1–18, 2014. <https://doi.org/10.1016/j.apnum.2013.08.006>
- [2] M. Sharifi, A. Abdi, M. Braś, and G. Hojjati. High order second derivative diagonally implicit multistage integration methods for ODEs. *Mathematical Modelling and Analysis*, **28** (1):53–70, 2023. <https://doi.org/10.3846/mma.2023.16102>

RECENT ADVANCES IN THE NUMERICAL SOLUTION OF ORDINARY AND FRACTIONAL DIFFERENTIAL EQUATIONS

LUIGI BRUGNANO

Università di Firenze

Dipartimento di Matematica e Informatica “U. Dini”, Viale Morgagni 67/A, 50134 Firenze, Italy

E-mail: luigi.brugnano@unifi.it

Recently, the class of Runge–Kutta methods named *Hamiltonian Boundary Value Methods (HBVMs)* has been introduced for the efficient numerical solution of Hamiltonian problems [5; 6]. This is a class of energy-conserving, low-rank R–K methods, which relies on the expansion of the vector field along the Legendre polynomial basis. In this respect, the methods can be also regarded as spectral methods in time and, in fact, they are able to reach a spectral accuracy [1]. Later on, this approach has been extended to cope with different kinds of differential problems [3], including fractional differential equations [7]. Concerning these latter equations, however, only recently, the basis for the expansion has been tailored for the problem at hand [2], thus resulting into a much more efficient procedure [4].

In this talk, the main facts concerning this approach will be recalled, along with the more recent application for solving fractional differential equations.

REFERENCES

- [1] P. Amodio, L. Brugnano, F. Iavernaro. Analysis of Spectral Hamiltonian Boundary Value Methods (SHBVMs) for the numerical solution of ODE problems. *Numer. Algorithms*, **83** (2020) 1489–1508. <https://doi.org/10.1007/s11075-019-00733-7>
- [2] L. Brugnano, K. Burrage, P. Burrage, F. Iavernaro. A spectrally accurate step-by-step method for the numerical solution of fractional differential equations. [arXiv:2310.10526](https://arxiv.org/abs/2310.10526) [math.NA], <https://doi.org/10.48550/arXiv.2310.10526>.
- [3] L. Brugnano, G. Frasca-Caccia, F. Iavernaro, V. Vespri. A new framework for polynomial approximation to differential equations. *Adv. Comput. Math.*, **48** (2022) 76. <https://doi.org/10.1007/s10444-022-09992-w>
- [4] L. Brugnano, G. Gurioli, F. Iavernaro. Numerical solution of FDE-IVPs by using Fractional HBVMs: the `fhbvm` code. [arXiv:2403.04916](https://arxiv.org/abs/2403.04916) [math.NA], <https://arxiv.org/abs/2403.04916>.
- [5] L. Brugnano, F. Iavernaro. *Line Integral Methods for Conservative Problems*. Chapman et Hall/CRC, Boca Raton, FL. 2016
- [6] L. Brugnano, F. Iavernaro. Line Integral Solution of Differential Problems. *Axioms*, **7(2)** (2018) 36. <https://doi.org/10.3390/axioms7020036>
- [7] L. Brugnano, F. Iavernaro. A general framework for solving differential equations. *Annali dell’Università di Ferrara*, **68** (2022) 243–258. <https://doi.org/10.1007/s11565-022-00409-6>

HYERS-ULAM STABILITY OF IMPLICIT VOLTERRA EQUATIONS ON TIME SCALES

ANDREJS REINFELDS¹ and SHRADDHA RAMANBHAI CHRISTIAN²

¹*Institute of Mathematics and Computer Science, University of Latvia*

Raiņa bulv. 29, Rīga LV-1459, Latvia

²*Institute of Applied mathematics, Riga Technical University*

Zunda Krastmala 10, Rīga LV-1048, Latvia

E-mail: reinf@latnet.lv, Shraddha-Ramanbhai.Christian@rtu.lv

Consider nonlinear k -th order Volterra integrodifferential equation on an arbitrary time scale \mathbb{T}

$$x^{\Delta^k}(t) = f\left(t, x(t), x^{\Delta}(t), \dots, x^{\Delta^k}(t), \int_{t_0}^t k(t, s, x(s), x^{\Delta}(s), \dots, x^{\Delta^k}(s)) \Delta s\right) \quad (1)$$

with initial conditions

$$x^{\Delta^i}(t_0) = x_i, \quad i = 0, 1, 2, \dots, k-1, \quad t_0, t \in I_{\mathbb{T}} = [t_0, +\infty) \cap \mathbb{T}.$$

We reduce equation (1) to implicit Volterra integral equation

$$z(t) = F\left(t, z(t), \int_{t_0}^t K(t, s, z(s)) \Delta s\right), \quad t_0, t \in I_{\mathbb{T}} = [a, +\infty) \cap \mathbb{T}, \quad (2)$$

where $z: I_{\mathbb{T}} \rightarrow \mathbb{R}^{n(k+1)}$ is the unknown function, $K: I_{\mathbb{T}} \times I_{\mathbb{T}} \times \mathbb{R}^{n(k+1)} \rightarrow \mathbb{R}^{n(k+1)}$ be rd-continuous in its first and second variable, $L: I_{\mathbb{T}} \rightarrow \mathbb{R}$ be rd-continuous, $\gamma > 1$, $\beta = L(s)\gamma$,

$$|K(t, s, z) - K(t, s, z')| \leq L(s)|z - z'|, \quad (z, z') \in \mathbb{R}^{n(k+1)}, \quad s < t,$$

$$|F(t, z, w) - F(t, z', w')| \leq M(|z - z'| + |w - w'|), \quad m = \sup_{t \in I_{\mathbb{T}}} \frac{1}{e_{\beta}(t, t_0)} \left| F(t, 0, \int_{t_0}^t K(t, s, 0) \Delta s \right| < \infty.$$

If $M(1 + 1/\gamma) < 1$, then the integral equation (2) has a unique solution $z \in C_{\beta}^k(I_{\mathbb{T}}; \mathbb{R}^{n(k+1)})$, where $C_{\beta}^k(I_{\mathbb{T}}; \mathbb{R}^{n(k+1)})$ be the Banach space of rd-continuous functions such that

$$\sup_{t \in I_{\mathbb{T}}} \frac{\max_{0 \leq i \leq k} |x^{\Delta^i}(t)|}{e_{\beta}(t, t_0)} < \infty.$$

We also prove that equation (1) is Hyers-Ulam stable.

REFERENCES

- [1] A. Reinfelds and S. Christian. Nonlinear Volterra integrodifferential equations from above on unbounded time scales. *Mathematics*, **11** (7): 1760, 2023.
- [2] I. Daniela and M. Daniela. Semi-Hyers-Ulam-Rassias stability of some Volterra integro-differential equations via Laplace transform. *Axioms*, **12** (3): 279, 2023.

LONG STANDING AND NEVER OLD QUESTION: WHO IS A BOSS (STABILITY OR APPROXIMATION)

RAIMONDAS ČIEGIS

Vilnius Gediminas technical university

Saulėtekio al. 11, LT-10223 Vilnius, Lithuania

E-mail: rc@vgtu.lt

Adaptive grids in space and time are used to fit the grid points to the dynamics of the solution and to minimize the approximation/truncation error.

At the same time uniform grids are used in many recent big data projects due to two very important properties: high order approximations can be constructed directly on uniform grids and the obtained structure of grids is well suited for parallel computing techniques. As a consequence different modifications and new discrete schemes are proposed which try to preserve the uniformity of the grids as close as possible. The well known NFL theorem predicts that some additional costs should be expected. In our talk we consider some new interesting discrete schemes and show that the stability analysis should be tailored to each source of approximation errors.

A linear hyperbolic problem is approximated by the following implicit symmetrical three-level scheme

$$\frac{U^{n+1} - 2U^n + U^{n-1}}{\tau^2} + \beta \frac{U^{n+1} - U^{n-1}}{2\tau} + A_h \frac{U^{n+1} + U^{n-1}}{2} = F^n, \quad (1)$$
$$U^0 = u_0, \quad U^1 = u_0 + \tau v_0.$$

In [1], this scheme is modified for the specific quasi-uniform time grid when at some grid points the lengths of grid steps are doubled or halved. The most valuable property of this scheme is that the approximation is still done on uniformly distributed grid points, thus basic advantages of such discrete schemes are preserved.

By making a full stability analysis of interpolation errors introduced by the proposed algorithm we prove that the cases of doubling and reducing twice the time steps lead to totally different error accumulation rates. Results of computational experiments are presented and they confirm the accuracy of theoretical error estimates.

We also investigate the difference in the stability of the backward Euler (BE) finite difference scheme and the discontinuous Galerkin (DG) finite element scheme when both schemes are used to solve one dimensional parabolic problem on dynamically shifted uniform space grids. The theoretical analysis proves that the accumulation of interpolation and projection approximation errors are quite different.

Finally we ask the question if adaptive grids are really required and are more efficient than uniform grids?

REFERENCES

- [1] P.N. Vabishchevich, Three-level schemes with double change in the time step. *Computational Mathematics and Mathematical Physics*, **63** (11), 1989-1995 (2023)

MODELLING OF BACTERIAL PLUME FORMATION IN A CIRCULAR CONTAINER

BOLESLOVAS DAPKŪNAS

Institute of Computer Science, Vilnius University

Didlaukio g. 47, LT-08303 Vilnius, Lithuania

E-mail: boleslovas.dapkunas@mif.vu.lt

Microorganisms such as bacteria *Escherichia coli* move around either towards attractants (e.g. food) or away from repellants (e.g. toxins). This process is called chemotaxis, and it allows populations of bacteria to self-organize and form patterns [1]. One of the most widely used methods to model chemotaxis is the Keller-Segel approach [2].

E. coli pattern formation can be modelled using a system of reaction-diffusion-chemotaxis equations, representing dynamics of bacteria, self-excreted chemoattractant, and oxygen [3]. In this work, the model was coupled with Navier-Stokes equations to improve the modelling of plume formation [4]. The dimensionless 2D in space model is governed by these equations:

$$\frac{\partial n}{\partial t} + (\vec{u} \cdot \vec{\nabla}) n = D_n \Delta n - \chi \nabla \cdot (n \nabla c) + \alpha n \left(1 - \frac{n}{o}\right) \quad (1)$$

$$\frac{\partial c}{\partial t} + (\vec{u} \cdot \vec{\nabla}) c = D_c \Delta c + \frac{n}{1 + \beta n} - c \quad (2)$$

$$\frac{\partial o}{\partial t} + (\vec{u} \cdot \vec{\nabla}) o = D_o \Delta o - \lambda n \quad (3)$$

$$\frac{\partial \omega}{\partial t} + (\vec{u} \cdot \vec{\nabla}) \omega = \nu \Delta \omega - \kappa \frac{\partial n}{\partial x} \quad (4)$$

$$\Delta \Psi = -\omega, \quad (x, y) \in (0, l) \times (0, h), \quad t > 0, \quad (5)$$

where $n(x, y, t)$, $c(x, y, t)$, and $o(x, y, t)$ represent cell density, chemoattractant, and oxygen respectively, $\omega(x, y, t)$ is the vorticity, and $\Psi(x, y, t)$ is the stream function. Together with appropriate initial and boundary conditions, the governing equations form a boundary-value problem, which was solved using finite difference technique.

The aim of this work is to investigate the effects of model parameters on bacteria pattern formation, with special emphasis on plume formation.

REFERENCES

- [1] E. O. Budrene and H. C. Berg. Dynamics of formation of symmetrical patterns by chemotactic bacteria. *Nature*, **376** (6535):49–53, 1995.
- [2] K. J. Painter. Mathematical models for chemotaxis and their applications in self-organisation phenomena. *Journal of Theoretical Biology*, **481** 162–182, 2019.
- [3] R. Šimkus, R. Baronas and Ž. Ledas. A multi-cellular network of metabolically active *E. coli* as a weak gel of living Janus particles. *Soft Matter*, **9** (17):4489–4500, 2013.
- [4] A. Chertock, K. Fellner, A. Kurganov, A. Lorz and P. A. Markowich. Sinking, merging and stationary plumes in a coupled chemotaxis-fluid model: a high-resolution numerical approach. *Journal of Fluid Mechanics*, **694** 155–190, 2012.

FAST NUMERICAL ALGORITHMS FOR THE EVALUATION OF RIEMANN-LIOUVILLE INTEGRALS

KAI DIETHELM

*Technical University of Applied Sciences Würzburg-Schweinfurt, Faculty of Applied Natural
Sciences and Humanities*

Ignaz-Schön-Str. 11, 97421 Schweinfurt, Germany

E-mail: kai.diethelm@thws.de

In many applications, it is necessary to compute numerical solutions of one or many ordinary differential equations of fractional order over many time steps. At the core of such numerical solvers, we typically find schemes for the approximate evaluation of Riemann-Liouville integrals. Due to the inherent memory properties of the associated differential and integral operators, such computations require a significant amount of computer memory and a relatively large runtime if straightforward discretization approaches are used. More sophisticated techniques, such as algorithms based on the FFT, can reduce the runtime significantly but not the memory requirements. Other concepts like, e.g., nested meshes reduce the memory demands but come with a significant administrative overhead. The goal of this talk is to present a class of algorithms that simultaneously address both the runtime and the memory footprint issues.

Specifically, our methods are based on so-called diffusive representations of the Riemann-Liouville integral operator. This allows us to handle the process memory in an implicit manner, thus immediately reducing the algorithm's required computer memory to an order of magnitude comparable to that of an algorithm for local problems. Many different analytic variants of such representations exist, and each of them allows multiple combinations of algorithms for its numerical evaluation. It is therefore necessary to identify special cases that admit a particularly accurate numerical approximation with a small number of arithmetic operations, thus additionally addressing the runtime matter in a satisfactory way.

The work described in this talk is performed within a joint project with Renu Chaudhary and Afshin Farhadi (THWS) and André Schmidt and Paul E. Haacker (Institute of Nonlinear Mechanics, Universität Stuttgart). This project is supported by the German Federal Ministry of Education and Research (BMBF) under Grant 05M22WHA.

CO AND CO₂ CONCENTRATION IN BIOMASS COMBUSTION PROCESS DEPENDING ON BIOMASS POROSITY


MARIS GUNARS DZENIS

University of Latvia

Jelgavas street 3, Riga, Latvia

E-mail: maris_gunars.dzenis@lu.lv

To simulate thermal decomposition of biomass we define every biomass as three organic compounds. Organic compound thermal decomposes in volatile part and carbon part [2]. Coal combustion are described with equation



Reactions are modeled using Arrhenius kinetics. To determine changes off biomass that microwave pretreatment gives we estimate these changes using porosity. Our objective is to determine how porosity changes in biomass affects CO and CO₂ concentration in mathematical model. Gases we model using the Darcy law, ideal gas law and mass balance equation [1]. Numerical solutions were found using finite difference scheme and finite volume method in program MatLab.

REFERENCES

- [1] U. Strautins, L. Leja, M. G. Dzenis. Some Network Models Related to Heat and Mass Transfer During Thermal Conversion Of Biomass. *Enginering for rural development*, **20** 1213 - 1218, 2021.
- [2] Goldšteins, L., Dzenis, M.G., Šints, V., Zake, M., Arshanitsa, A.. Microwave Pre-Treatment and Blending of Biomass Pellets for Sustainable Use of Local Energy Resources in Energy Production. *Energies*, **15(3)** 755, 2022.

A GLOBAL APPROXIMATION METHOD FOR SECOND-KIND VOLTERRA-FREDHOLM EQUATIONS

LUISA FERMO, DOMENICO MEZZANOTTE, and DONATELLA OCCORSIO

Department of Mathematics and Computer Science, University of Cagliari

Via Ospedale 72, 09124 Cagliari, Italy

E-mail: fermo@unica.it

Department of Mathematics and Computer Science, University of Basilicata

Via dell'Ateneo Lucano 10, 85100 Potenza, Italy

E-mail: donatella.occorsio@unibas.it

Department of Mathematics "Giuseppe Peano", University of Torino

Via Verdi 8, 10124 Torino, Italy

E-mail: domenico.mezzanotte@unito.it

This talk deals with the numerical treatment of the following mixed Volterra-Fredholm integral equation

$$(I + \mu VK)f = g,$$

where $\mu \in \mathbb{R} \setminus \{0\}$, f is the unknown function, g is a given right-hand side, I is the identity operator, V is the Volterra operator given by

$$(Vf)(y) = \int_{-1}^y h(x, y)f(x)(y-x)^\rho(1+x)^\sigma dx, \quad \rho, \sigma > -1, \quad y \in [-1, 1],$$

with h an assigned kernel, and K is the Fredholm operator defined as

$$(Kf)(y) = \int_{-1}^1 k(x, y)f(x)(1-x)^\alpha(1+x)^\beta dx, \quad \alpha, \beta > -1, \quad y \in [-1, 1],$$

with k a known kernel.

A global approximation method of Nyström type based on a mixed Gauss-product cubature formula is developed to approximate the solution f in suitable weighted spaces equipped with the uniform norm. Stability and convergence results are discussed and numerical tests are given to show the good performance of the method.

INTEGRATING FRACTIONAL DIFFERENTIAL EQUATIONS WITH NEURAL NETWORKS FOR DYNAMICAL SYSTEMS MODELLING

CECÍLIA COELHO¹, M. FERNANDA P. COSTA¹, LUÍS L. FERRÁS^{1,2}

¹*Centre of Mathematics (CMAT), University of Minho*

²*Department of Mechanical Engineering (Section of Mathematics), FEUP - University of Porto*
Portugal

E-mail: cmartins@cmat.uminho.pt, mfc@math.uminho.pt, lferras@fe.up.pt

This study integrates Fractional Differential Equations (FDE) with Neural Networks for the modelling of dynamical systems. Inspired by concepts like Neural Ordinary Differential Equations (Neural ODE) [1] and fractional calculus within neural systems [2], we propose the Neural FDE [3]. In this neural network architecture, parameterised by θ (representing weights and biases), the objective is to determine the function f_θ of a Fractional Differential Equation of order α (equation (1)), ensuring that the solution to (1) accurately fits provided data (a time series), by adjusting the weights and biases within f_θ [3],

$${}_0^C D_t^\alpha \mathbf{h}(t) = \mathbf{f}_\theta(t, \mathbf{h}(t)) \text{ with } \mathbf{h}(t_0) = \mathbf{h}_0, \quad \alpha = \alpha_\phi. \quad (1)$$

Here, ${}_0^C D_t^\alpha \mathbf{h}(t)$ denotes the Caputo fractional derivative [4], where $\mathbf{h}(t)$ represents the state of the dynamical system at time t , and $\mathbf{h}(t_0) = \mathbf{h}_0$ is the initial condition. As this model introduces an additional parameter (α) compared to the Neural ODE, the order of the Fractional Differential Equation, α , is also learned by another neural network, α_ϕ , with parameters ϕ .

This hybrid fractional differential equation can then be used for making future predictions or predicting missing data. The solution of (1) is obtained numerically, with the *values* of $f_\theta(t, \mathbf{h}(t))$ provided by the neural network.

We limit our consideration to $\alpha \in (0, 1)$, as it finds applicability in various scenarios and simplifies the formulation.

REFERENCES

- [1] R.T. Chen, Y. Rubanova, J. Bettencourt and D.K. Duvenaud. Neural ordinary differential equations. In: *Proc. of the Intern. Conference Advances in Neural Information Processing Systems 31 (NeurIPS 2018), Montréal, Canada, 2018*, Navier-Stokes Equations and Related Nonlinear Problems, S. Bengio, H. Wallach, H. Larochelle, K. Grauman, N. Cesa-Bianchi and R. Garnett (Eds.), 2018, 1 – 13.
- [2] B.N. Lundstrom, M.H. Higgs, W.J. Spain and A.L. Fairhall. Fractional differentiation by neocortical pyramidal neurons. *Nature Neuroscience*, **11** 335–1342, 2008.
- [3] C. Coelho, M.F.P. Costa and L.L. Ferrás. Neural Fractional Differential Equations. <https://doi.org/10.48550/arXiv.2403.02737>, .
- [4] K. Diethelm. *The Analysis of Fractional Differential Equations: An Application - Oriented Exposition Using Differential Operators of Caputo Type*. Springer-Verlag, Berlin Heidelberg, 2010.

MODELLING AND SIMULATION OF SYSTEMS WITH COMPLEX CHARACTERISTICS: CERTAINTIES, UNCERTAINTIES AND COMPROMISES

NEVILLE J FORD

University of Chester, UK

Parkgate Road, Chester, CH1 4BJ, UK

E-mail: njford@chester.ac.uk

In this talk we discuss the link between real world systems and the development of appropriate model equations. Our focus is on problems that involve a history, delay or after-effect, and we begin with a discussion of how these phenomena can be reflected in relatively simple model equations, applying classical results from [2] and [4]. This leads to a discussion of characteristic values and eigenfunctions and an understanding of the characteristics of the solution space and the limitations imposed by modelling choices. Here we apply more recent insights from [6].

We discuss approaches to model selection and parameter estimation and we consider how to match the purpose of the model with its formulation. We give examples that show how different modelling paradigms can be necessary to develop effective models in different circumstances (see, for example, [3] for further details on this theme).

We use the recent COVID-19 pandemic as a case study where model equations were widely used but poorly understood (see [5] and [1]). We give examples that show why the conclusions drawn from the model may be quite unhelpful to the application and we provide some ideas about how to avoid these problems arising in future modelling projects.

REFERENCES

- [1] Adiga, A., Dubhashi, D., Lewis, B., Marathe, M., Venkatramanan, S., & Vullikanti, Mathematical Models for COVID-19 Pandemic: A Comparative Analysis, *Computer Methods and Programs in Biomedicine*, 209, 106301, 2021.
- [2] Bellman, R. & Cooke, K.L., *Differential-Difference Equations*, Academic Press, 1963.
- [3] Bocharov, G.A., & Rihan, F.A., Numerical modelling in biosciences using delay differential equations, *Journal CAM*, 125, p183-199, 2000.
- [4] Diekmann, O., van Gils, S.A., Verduyn Lunel, S.M. & Walther, H.-O., *Delay Equations, Functional, Complex and Nonlinear Analysis*, Springer-Verlag, 1995.
- [5] Shankar S., Mohakuda S.S., Kumar A., Nazneen P.S., Yadav A.K., Chatterjee K., & Chatterjee K., Systematic review of predictive mathematical models of COVID-19 epidemic. *Med J Armed Forces India*. 2021 Jul., 77(Suppl 2), S385-S392, 2021.
- [6] Yi, S., Nelson, P.W., & Ulsoy, A.G., *Time-delay systems: Analysis and Control using the Lambert W function*, World Scientific, 2010.

A NONLINEAR BACKWARD DIFFUSION ALGORITHM FOR IMAGE RESTORATION

LORELLA FATONE

Università di Camerino

Via Madonna delle Carceri 9, Camerino, 62032, Italy

E-mail: `lorella.fatone@unicam.it`

DANIELE FUNARO

Università di Modena e Reggio Emilia

Via Campi 103, Modena, 41125, Italy

E-mail: `daniele.funaro@unimore.it`

We discuss the recent image processing methodology introduced in [2] in the context of nonlinear diffusive PDEs (see also [1; 3; 4; 5; 7]). A corrupted image u is represented in a continuous framework and used as initial guess for the evolution equation:

$$\frac{\partial u}{\partial t} = \sqrt{\frac{\|\nabla u\|^2}{1 + \|\nabla u\|^2}} \Delta u, \quad (1)$$

where ∇ and Δ denote the gradient and the Laplace operator, respectively. Equation (1) is defined on a rectangle and Neumann conditions are imposed on the boundaries. Contrary to the classical approaches, descending for example from the pioneering work of Perona & Malik [6], the nonlinear diffusion coefficient depends locally on the magnitude of $\|\nabla u\|$ and grows proportionally to this quantity. An ad-hoc discretization scheme uses centered finite-differences scheme for the space variable and it is implemented on two interlaced grids. Euler's scheme is applied for time discretization going backwards in time. The aim is to smooth out the regions of almost uniform shade and emphasize discontinuities. The proposed methodology is quite efficient as satisfactory results are obtained after only one time step of appropriate size. The algorithm is well-suited for edge detection and segmentation problems. We show some applications in the case of images affected by blur and noise.

REFERENCES

- [1] S. Bonettini, F. Porta, V. Ruggiero and L. Zanni: Variable metric techniques for forward-backward methods in imaging, *J. Comput. Appl. Math.* 385:1–30, 2021.
- [2] L. Fatone and D. Funaro, High-order discretization of backward anisotropic diffusion and application to image processing, *Ann. Università di Ferrara* 68(2):295–310, 2022.
- [3] M. Felsberg, On the relation between anisotropic diffusion and iterated adaptive filtering, *LNCS*, 436–445, 2008.
- [4] M. Grasmair and F. Lenzen, Anisotropic total variation filtering, *Appl. Math. Optim.* 62:323–339, 2010.
- [5] M. Pragliola, L. Calatroni, A. Lanza and F. Sgallari, On and beyond total variation in imaging: the role of space variance, *SIAM Rev.* 65(3):601–685, 2023.
- [6] P. Perona and J. Malik, Scale space and edge detection using anisotropic diffusion, *IEEE Trans. on Pattern Anal. and Machine Intelligence* 12(7):629–639, 1990.
- [7] C. Solomon and T. Breckon, *Fundamentals of Digital Image Processing: A Practical Approach With Examples in Matlab*, John Wiley & Sons, 2010.

ON DISCRETE SHIFTS OF THE MELLIN TRANSFORM OF THE RIEMMAN ZETA-FUNCTION

VIRGINIJA GARBALIAUSKIENĖ^{1,2}, ANTANAS LAURINČIKAS³ and DARIUS
ŠIAUČIŪNAS²

¹*Šiaulių valstybinė kolegija*

Aušros av. 40, LT-76241 Šiauliai, Lithuania

²*Šiauliai Academy, Vilnius University*

Vytauto street 84, LT-76352 Šiauliai, Lithuania

E-mail: virginija.garbaliuskiene@sa.vu.lt, darius.siauciunas@sa.vu.lt

³*Faculty of Mathematics and informatics, Vilnius University*

Naugarduko street 24, LT-03225 Vilnius, Lithuania

E-mail: antanas.laurincikas@mif.vu.lt

We consider the Mellin transform $\mathcal{Z}(s)$, $s = \sigma + it$, of the square of the Riemann zeta-function $\zeta(s)$, i.e.,

$$\mathcal{Z}(s) = \int_1^\infty |\zeta(1/2 + ix)|^2 x^{-s} dx,$$

and the approximation of analytic functions by discrete shifts $\mathcal{Z}(s + ikh)$ with $h > 0$ and $k \in \mathbb{N}_0 = \mathbb{N} \cup \{0\}$. Let $D = \{s \in \mathbb{C} : 1/2 < \sigma < 1\}$, and $H(D)$ denote the space of analytic functions on D equipped with the topology of uniform convergence on compacta.

In the report, we will discuss the following theorem, for details, see [1].

THEOREM 1. *For $h > 0$, there exists a non-empty closed set F_h such that, for every compact set $K \subset D$, $f(s) \in F_h$ and $\varepsilon > 0$,*

$$\liminf_{T \rightarrow \infty} \frac{1}{N+1} \# \left\{ 0 \leq k \leq N : \sup_{s \in K} |\mathcal{Z}(s + ikh) - f(s)| < \varepsilon \right\}.$$

Moreover, the limit

$$\lim_{T \rightarrow \infty} \frac{1}{N+1} \# \left\{ 0 \leq k \leq N : \sup_{s \in K} |\mathcal{Z}(s + ikh) - f(s)| < \varepsilon \right\}$$

exists and is positive for all but at most countably many $\varepsilon > 0$.

REFERENCES

- [1] V. Garbaliuskienė, A. Laurinčikas and D. Šiaučiūnas. On the discrete approximation by Mellin transform of the Riemann zeta-function. *Mathematics*, **11** (10):article no. 2315, 2023.

NUMERICAL APPROACHES TO SOLVE AN INVERSE PROBLEM FOR A FRACTIONAL DIFFUSION-WAVE EQUATION WITH MULTI-TERM FRACTIONAL POWERS OF MINUS LAPLACIAN

HANY GERGES, JAAN JANNO

Department of Cybernetics, School of Science, Tallinn University of Technology
Ehitajate tee 5, 19086 Tallinn, Estonia
E-mail: hagerg@taltech.ee

Two numerical approaches designed using MATLAB to recover the unknowns of the following fractional diffusion-wave equation with multi-term fractional powers of minus Laplacian

$$D_t^\alpha u(x, t) = \sum_{j=1}^n b_j (-\Delta)^{\beta_j} u(x, t) + f_0(x, t), \quad (1)$$

are considered. The first approach is based on the minimization of the error of the Laplace transform of the observation function minus the Laplace transform of the observation function with the unknowns as variables and the second approach uses location of poles of Laplace transform of observation function. Finally, a comparison of the two approaches is presented.

Uniqueness for the solution is proved theoretically, also by means of location of poles.

REFERENCES

- [1] K.Liao, T.Wei. Identifying a fractional order and a space source term in a time-fractional diffusion-wave equation simultaneously. *Inverse Problems*, **35** 115002, 2019.
- [2] J. Janno, N. Kinash. Reconstruction of an order of derivative and a source term in a fractional diffusion equation from final measurements. *Inverse Problems*, **34** 025007, 2018.
- [3] A.A. Kilbas, H.M. Srivastava, J.J. Trujillo. Theory and Applications of Fractional Differential Equations *Elsevier, Amsterdam*. 2006, .

JOINT MIXED LIMIT THEOREM FOR EPSTEIN AND HURWITZ ZETA-FUNCTIONS

HANY GERGES, RENATA MACAITIENĖ *

Institute of Mathematics, Faculty of Mathematics and Informatics, Vilnius University

Naugarduko str. 24, LT-03225 Vilnius, Lithuania

E-mail: hany.gerges@mif.stud.vu.lt

In the talk, we will present some known results on the joint asymptotic behaviour of zeta-functions having and having no Euler's product over prime numbers. Moreover, we will discuss a new result for the Epstein and Hurwitz zeta-functions, more precisely, we will prove the joint probabilistic limit theorem in terms of the weak convergence of probability measures on \mathbb{C}^2 defined by means of these zeta-functions.

We recall the definitions of the mentioned functions.

The Epstein zeta-function $\zeta(s; Q)$ was introduced by P. Epstein in [1]. Let Q be a positive definite quadratic $n \times n$ matrix and $Q[\underline{x}] = \underline{x}^T Q \underline{x}$ for $\underline{x} \in \mathbb{Z}^n$. The Epstein zeta-function $\zeta(s; Q)$, $s = \sigma + it$, is defined, for $\sigma > \frac{n}{2}$, by the series $\zeta(s; Q) = \sum_{\underline{x} \in \mathbb{Z}^n \setminus \{0\}} (Q[\underline{x}])^{-s}$, and can be continued analytically to the whole complex plane, except for a simple pole at the point $s = \frac{n}{2}$ with residue $\frac{\pi^{n/2}}{\Gamma(n/2)\sqrt{\det Q}}$.

The Hurwitz zeta-function $\zeta(s, \alpha)$ was introduced in [2]. The function $\zeta(s, \alpha)$ with a fixed parameter α , $0 < \alpha \leq 1$, is defined, for $\sigma > 1$, by $\zeta(s, \alpha) = \sum_{m=1}^{\infty} (m + \alpha)^{-s}$, and has analytic continuation to the whole complex plane, except for a simple pole at the point $s = 1$ with residue 1.

REFERENCES

- [1] P. Epstein. Zur Theorie allgemeiner Zetafunktionen. *Math. Ann.*, **56**: 615–644, 1903.
- [2] A. Hurwitz. Einige Eigenschaften der Dirichlet'schen Functionen $F(x) = \sum \left(\frac{D}{n}\right) \frac{1}{n^s}$, die bei der Bestimmung der Klassenzahlen binärer quadratischer Formen auftreten. *Schlömilch Z.*, **27**: 86–102, 1882.

*The research of R. Macaitienė is funded by the Research Council of Lithuania, agreement No S-MIP-22-81.

FRACTIONAL CALCULUS OF ZETA FUNCTIONS

EMANUEL GUARIGLIA

Wenzhou-Kean University

88 Daxue Rd, 325060 Wenzhou, China

E-mail: eguarigl@kean.edu

This talk outlines fractional calculus of zeta functions. The main results are based on a complex generalization of the Grünwald-Letnikov fractional derivative. The functional equations can be rewritten in a simplified form. Thus, we reduce their computational cost. Moreover, for the case of the Riemann ζ function we propose a quasisymmetric form of the corresponding functional equation. The second part of this talk deals with the link with distribution of prime numbers. In particular, we show some results on free-zero regions for this class of functions. Finally, we discuss the representation of these functions in terms of Euler products.

REFERENCES

- [1] Apostol, T.M. Formulas for higher derivatives of the Riemann zeta function. *Math. Comp.* 1985;44(169):223–232.
- [2] Choudhury, B.K. The Riemann zeta-function and its derivatives. *Proc. R. Soc. Lond. A* 1995;450(1940):477–499.
- [3] Guariglia, E. Fractional calculus of the Lerch zeta function. *Mediterr J Math.* 2022;19(3):Art. ID 109.
- [4] Guariglia E. Fractional calculus, zeta functions and Shannon entropy. *Open Math.* 2021;19(1):87–100.
- [5] Guariglia E. Riemann zeta fractional derivative – functional equation and link with primes. *Adv. Difference Equ.* 2019;2019(1):Art. ID 261.
- [6] Li, C., Dao, X., Guo, P. Fractional derivatives in complex planes. *Nonlinear Anal.* 2009;71(5-6):1857–1869.
- [7] Ortigueira, M.D., Coito, F.J. From differences to derivatives. *Fract. Calc. Appl. Anal.* 2004;7(4):459–471.

LOCAL ERROR ESTIMATION FOR IMPLICIT-EXPLICIT GENERAL LINEAR METHODS

ZDZISŁAW JACKIEWICZ

Arizona State University

1151 S Forest Ave, Tempe, AZ, United States

E-mail: jackiewicz@asu.edu

We investigate implicit-explicit methods for differential systems with stiff and non-stiff parts. Stage order and order conditions are formulated and estimation of local discretization errors in fixed and variable stepsize environments is discussed. We also describe the construction of such methods with desirable accuracy and stability properties. This is a joint work with Angela Cardone, University of Salerno, and Giuseppe Izzo, University of Naples.

APPROXIMATION OF ANALYTIC FUNCTIONS BY SHIFTS OF ABSOLUTELY CONVERGENT DIRICHLET SERIES RELATED TO PERIODIC ZETA-FUNCTIONS

MINDAUGAS JASAS

Faculty of Mathematics and Informatics, Vilnius University

Naugarduko street 24, LT-03225 Vilnius, Lithuania

E-mail: mindaugas.jasas@mif.stud.vu.lt

Let $\mathbf{a} = \{a_m : m \in \mathbb{N}\}$ and $\mathbf{b} = \{b_m : m \in \mathbb{N}_0 = \mathbb{N} \cup \{0\}\}$ be two periodic sequences of complex numbers, $0 < \alpha \leq 1$ and $\theta > \frac{1}{2}$ are fixed parameters, $v_u(m) = \exp\left\{-\left(\frac{m}{u}\right)^\theta\right\}$, $m \in \mathbb{N}$, and $v_u(m, \alpha) = \exp\left\{-\left(\frac{m+\alpha}{u}\right)^\theta\right\}$, $m \in \mathbb{N}_0$, with $u > 0$. Let $s = \sigma + it$ denote a complex variable. We consider the Dirichlet series

$$\zeta_u(s; \mathbf{a}) = \sum_{m=1}^{\infty} \frac{a_m v_u(m)}{m^s} \quad \text{and} \quad \zeta_u(s, \alpha; \mathbf{b}) = \sum_{m=0}^{\infty} \frac{b_m v_u(m, \alpha)}{(m + \alpha)^s}$$

which are absolutely convergent in any half-plane $\sigma > \sigma_0$ with finite σ_0 . We obtain the following results:

1. Approximation of a class of analytic functions by continuous shifts
 $\zeta_{u_T}(s + i\tau; \mathbf{a})$, $\tau \in \mathbb{R}$, with multiplicative sequence \mathbf{a} , and $u_T \rightarrow \infty$ and $u_T \ll T^2$.
2. Approximation of a class of analytic functions by discrete shifts
 $\zeta_{u_N}(s + ikh; \mathbf{a})$, $h > 0$, $k \in \mathbb{N}_0$, with multiplicative sequence \mathbf{a} , and $u_N \rightarrow \infty$ and $u_N \ll N^2$.
3. Joint approximation of a class of pairs of analytic functions by continuous shifts
 $(\zeta_{u_T}(s + i\tau; \mathbf{a}), \zeta_{u_T}(s + i\tau, \alpha; \mathbf{b}))$, $\tau \in \mathbb{R}$, with multiplicative sequence \mathbf{a} , and $u_T \rightarrow \infty$ and $u_T \ll T^2$.
4. Joint approximation of a class of pairs of analytic functions by discrete shifts
 $(\zeta_{u_N}(s + ikh_1; \mathbf{a}), \zeta_{u_N}(s + ikh_2, \alpha; \mathbf{b}))$, $h_1 > 0$, $h_2 > 0$, $k \in \mathbb{N}_0$, with multiplicative sequence \mathbf{a} , and $u_N \rightarrow \infty$ and $u_N \ll N^2$.

More precisely, let K_1, K_2 be compact sets of the strip $D = \{s \in \mathbb{C} : \frac{1}{2} < \sigma < 1\}$ with connected complements, $f_1(s)$ continuous, nonvanishing and $f_2(s)$ continuous functions on K_1 and K_2 , and analytic in interior of K_1, K_2 , respectively.

Suppose that the set $\{(h_1 \log p : p \in \mathbb{P}), (h_2 \log(m + \alpha) : m \in \mathbb{N}_0), 2\pi\}$ is linearly independent over the field of rational numbers. Then the limit

$$\lim_{N \rightarrow \infty} \frac{1}{N+1} \#\left\{0 \leq k \leq N : \sup_{s \in K_1} |\zeta_{u_N}(s + ikh_1; \mathbf{a}) - f_1(s)| < \varepsilon_1, \right. \\ \left. \sup_{s \in K_2} |\zeta_{u_N}(s + ikh_2, \alpha; \mathbf{b}) - f_2(s)| < \varepsilon_2 \right\}$$

exists and is positive for all but at most countably many $\varepsilon_1 > 0$ and $\varepsilon_2 > 0$. The above results form PhD thesis of the author.

REGULARITY RESULTS FOR VERY WEAK TIME-PERIODIC POISEUILLE-TYPE SOLUTION WITH MINIMALLY REGULAR FLOW RATE

RITA JUODAGALVYTĖ

Faculty of Mathematics and Informatics, Vilnius University

Naugarduko street 24, LT-03225, Vilnius, Lithuania

E-mail: rita.juodagalvyte@mif.vu.lt

In this presentation, we provide a summary of our results concerning the time-periodic very weak solutions of the heat equation with a non-local additional condition of the prescribed flux

$$F(t) = \int_{\sigma} U(x, t) dx$$

and their regularity properties. Specifically, we present an example of a function $F(t)$ that belongs to $L^2(-\pi, \pi)$ but $F \notin W^{\beta, 2}(-\pi, \pi)$ for $0 < \beta < 1$. The very weak solution corresponding to such flux F , has the regularity as stated in the definition (see [1],[2]) and it is not better. Furthermore, we discuss results that suggest a correlation between the improvement of solutions regularity and the increase in regularity of the given function F .

REFERENCES

- [1] K. Pileckas, R. Čiegis. Existence of nonstationary Poiseuille type solutions under minimal regularity assumptions. *Z. Angew. Math. Phys.*, **71**, 192, 2020. <https://doi.org/10.1007/s00033-020-01422-5>
- [2] K. Kaulakytė, N. Kozulinas and K. Pileckas. Time-periodic Poiseuille-type solution with minimally regular flow rate. *Nonlinear analysis: modelling and control*, **26**, 5, 947–968, 2021. <https://doi.org/10.15388/namc.2021.26.24502>

NOTES ON SIMULTANEOUS APPROXIMATION BY THE CLASS OF ZETA-FUNCTIONS

ROMA KAČINSKAITĒ

Vilnius University

Naugarduko str. 24, LT-03225 Vilnius, Lithuania

E-mail: roma.kacinskaite@mif.vu.lt

The theory of universality focuses on the approximation of analytic functions by the shifts of certain zeta-functions. In a much more complicated situation, we touch on studying approximation by the wide classes of zeta-functions. One such class is the so-called Selberg-Steuding class $\tilde{\mathcal{S}}$, which was introduced by A. Selberg in [3] and modified by J. Steuding in [4], and is defined by Dirichlet series $L(s)$ satisfying certain axioms.

The first approximation result related to the class $\tilde{\mathcal{S}}$ was obtained by J. Steuding in [4] and later improved by H. Nagoshi and J. Steuding in [2]. R. Kačinskaitė, A. Laurinčikas and B. Žemaitienė have obtained [1] the joint universality theorem for L -functions belonging to the Selberg-Steuding class $\tilde{\mathcal{S}}$. More precisely, we have shown a result about simultaneous approximation of a collection of analytic functions $(f_1(s), \dots, f_r(s))$ in the strip $\{s \in \mathbb{C} : \sigma_L < \sigma < 1\}$ by a collection of shifts $(L(s + ia_1\tau), \dots, L(s + ia_r\tau))$, $L(s) \in \tilde{\mathcal{S}}$, where $\sigma_L > \frac{1}{2}$ is a number depending on L , and real algebraic numbers a_1, \dots, a_r are linearly independent over the field of rational numbers \mathbb{Q} .

In the talk, we discuss two results on the investigation of joint universality property for $L(s) \in \tilde{\mathcal{S}}$, i.e., we present the joint functional independence and denseness for the above-mentioned collection.

REFERENCES

- [1] R. Kačinskaitė, A. Laurinčikas, and B. Žemaitienė. Joint universality in the Selberg-Steuding class. *Mathematics*, **11** : 737, 2023.
- [2] H. Nagoshi, J. Steuding. Universality for L -functions in the Selberg class. *Lith. Math. J.*, **50** 3: 293–311, 2010.
- [3] A. Selberg. Old and new conjectures and results about a class of Dirichlet series. In: *Proc. of the Amalfi Conf. on Analytic Number Theory, Maiori, Amalfi, Italy, 1989*, E. Bombieri et al. (eds.), University di Salerno, Salerno, 1992, 367–385.
- [4] J. Steuding. *Value Distribution of L-Functions*. Springer, Berlin/Heidelberg, New York, 2007.

STALLING IN QUEUING SYSTEMS WITH HETEROGENEOUS CHANNELS

LIUDVIKAS KAKLAUSKAS

Faculty of Business and Technologies, Šiauliai State Higher Education Institution

Aušros av. 40, LT-76241 Šiauliai, Lithuania

E-mail: `l.kaklauskas@svako.lt`

In the talk, a model of stalling in the queueing system (QS) with any number of different capacities heterogeneous servers will be discussed and some new results published in [1] will be presented. The model of stalling in QSs with two heterogeneous servers has been considered in [2], where the explicit probabilities of steady states were derived.

We will discuss the optimization of a stalling buffer as well, and we will show that stalling helps us to solve the slow server problem under an appropriate choice of stalling buffer size, making the slow servers usable under various values of system load. Moreover, some applications of the developed model in heterogeneous server clusters and in work productivity modelling for forest harvesting applications will be presented.

REFERENCES

- [1] L. Sakalauskas, L. Kaklauskas and R. Macaitienė. Stalling in queueing systems with heterogeneous channels. *Applied Science*, **14** (2), art. no. 773: 1–16, 2024.
- [2] L. Kaklauskas, L. Sakalauskas and V. Denisovas. Stalling for solving slow server problem. *RAIRO Oper. Res*, **53** 1097–1107, 2019.

A NONLINEAR MATHEMATICAL MODEL OF THE COURSE AND THERAPY OF RHEUMATOID ARTHRITIS AND ITS REALIZATION

ZVIAD KALICHAVA

Niko Muskhelishvili Institute of Computational Mathematics of the Georgian Technical University

4 Grigol Peradze Str., Tbilisi 0159, Georgia

E-mail: zviadi.kalichava@gmail.com

KAKHABER ODISHARIA, VLADIMIR ODISHARIA

Ivane Javakishvili Tbilisi State University

13 University Str., Tbilisi 0186, Georgia

E-mail: kodisharia@gmail.com; vladimer.odisharia@tsu.ge

NONA JANIKASHVILI

Tbilisi State Medical University

33 Vazha Pshavela Ave., Tbilisi 0186, Georgia

E-mail: njanikashvili@tsmu.edu

Mathematical models of immune mediated disorders provide an analytic platform in which we can address specific questions concerning disease immune dynamics to dictate the choice of treatment. Rheumatoid arthritis is a systemic autoimmune disease characterized by the joint inflammation and the cartilage destruction. Autoreactive B lymphocytes represent the integral elements of the pathophysiology of rheumatoid arthritis. Immune balance between the effector and the regulatory T cell subsets guide the autoreactive B cell fate and play a cardinal role in disease severity. Using non-linear differential equations, we developed a novel mathematical model that describes the immunopathogenesis of rheumatoid arthritis [1; 2]. The model explores the functional dynamics of cartilage destruction during disease progression, in which a system of differential equations depicts the interactions between autoreactive B lymphocytes and T helper cells. As the further task, we present here the refined model of the disease course in which the immunomodulatory effect of IL-6, – a molecule that drives the cross-talk of pro-inflammatory and regulatory subsets of T lymphocytes, is explained. IL-6 targeting is also taken into consideration in the disease treatment model, in which the modalities of treatment with methotrexate and tocilizumab in a separate or combined scheme are addressed. For such treatment model, the corresponding Cauchy problem is posed and its solution is found. In conclusion, we propose a novel mathematical model that best describes the readouts of the course and treatment outcomes of rheumatoid arthritis and, therefore, may take a rapid pace towards its implementation in biomedical and clinical research.

Acknowledgements. Grant support: Shota Rustaveli National Science Foundation of Georgia (grant # STEM 22-360).

REFERENCES

- [1] K. Odisharia, V. Odisharia, P. Tsereteli and N. Janikashvili. On the mathematical model of drug treatment of rheumatoid arthritis. *Mathematics, Informatics, and their Applications in Natural Sciences and Engineering*, **10** 161–168, 2019.
- [2] P. Tsereteli, V. Odisharia and N. Janikashvili. Mathematical modeling of rheumatoid arthritis and its treatment. *Computer Sciences and Telecommunications*, **61** (1):19–31, 2022.

LEARNING HAMILTONIAN DYNAMICS WITH STRUCTURE-PRESERVING NEURAL NETWORKS AND DIMENSIONALITY REDUCTION

DĀVIS KALVĀNS, JĀNIS BAJĀRS

Faculty of Physics, Mathematics and Optometry, University of Latvia

Jelgavas street 3, Riga LV-1004, Latvia

E-mail: davis.kalvans@lu.lv, janis.bajars@lu.lv

Data-driven approaches employing structure-preserving algorithms, such as symplectic neural networks known as *SympNets* [1], have gained recognition for learning Hamiltonian systems' dynamics. Despite their promising results, the challenge of high dimensionality persists. In this work, we investigate dimensionality reduction techniques to model Hamiltonian systems effectively in lower-dimensional subspaces, thereby reducing training times while preserving prediction accuracy.

We focus on learning nonlinear localized wave solutions in a one-dimensional crystal lattice model, as the lattice can be of an arbitrary dimension and, as we show, the problem lends well to being modeled in a lower-dimensional subspace. That is done by employing dimensionality reduction techniques, such as the non-symplectic *Proper Orthogonal Decomposition* (POD) [2] and geometric structure-preserving *Proper Symplectic Decomposition* (PSD) [3].

Moreover, we extend our previous work [4] by imposing symplectic neural networks architecture's map to be symmetric, i.e., equal to its adjoint map, as proposed in [5], a characteristic inherent to Hamiltonian system flows. Our results demonstrate that incorporation of the additional flows' property in symplectic neural networks together with structure-preserving dimensionality reduction enhances model predictions and their *valid prediction time* (VPT) even further.

REFERENCES

- [1] Pengzhan Jin, Zhen Zhang, Aiqing Zhu, Yifa Tang and George Em Karniadakis. *SympNets: Intrinsic structure-preserving symplectic networks for identifying Hamiltonian systems*. *Neural Networks*, **132** 166-179, 2020.
- [2] Steven Brunton and José Kutz. *Data-Driven Science and Engineering: Machine Learning, Dynamical Systems, and Control*. Cambridge: Cambridge University Press, 2019.
- [3] Liqian Peng and Kamran Mohseni. *Symplectic model reduction of Hamiltonian systems*. *SIAM Journal on Scientific Computing*, **38.1** A1-A27, 2016.
- [4] Jānis Bajārs and Dāvis Kalvāns. *Structure-preserving dimensionality reduction for learning Hamiltonian dynamics*. *Submitted*, 2023.
- [5] Jānis Bajārs. *Locally-symplectic neural networks for learning volume-preserving dynamics*. *Journal of Computational Physics*, **476** 111911, 2023.

ON CONVERGENCE CONDITIONS IN SELF-REGULARIZATION OF ILL-POSED PROBLEMS BY PROJECTION METHODS

URVE KANGRO, UNO HÄMARIK

Institute of Mathematics and Statistics, University of Tartu

Narva mnt 18, Tartu, 51009, Estonia

E-mail: urve.kangro@ut.ee, uno.hamarik@ut.ee

We consider an operator equation

$$Au = f, \quad f \in R(A) \neq \overline{R(A)},$$

where $A \in L(E, F)$ is the linear continuous operator between Banach spaces E and F . Instead of the exact right-hand side f we have only an approximation $f^\delta \in F$ satisfying condition $\|f - f^\delta\| \leq \delta$ with known δ . We consider projection methods. Let $E_n \subset E$, $Z_n \subset F^*$, $n \in \mathbb{N}$, be finite-dimensional nontrivial subspaces which have the role of approximating the spaces E and F^* , respectively. Let Q_n be the linear operator defined by

$$Q_n : F \rightarrow Z_n^* \quad \forall g \in F, z_n \in Z_n : \langle Q_n g, z_n \rangle_{Z_n^*, Z_n} = \langle z_n, g \rangle_{F^*, F}. \quad (1)$$

Then the finite-dimensional approximation $u_n \in E_n$ to the solution u_* can be found from the equation

$$Q_n A u_n = Q_n f^\delta. \quad (2)$$

Under conditions $\dim(E_n) = \dim(Z_n)$, $\mathcal{N}(Q_n A) \cap E_n = \{0\}$ the operator $A_n := Q_n A|_{E_n} : E_n \rightarrow Z_n^*$ has an inverse. In case of exact data ($\delta = 0$) the convergence $\|u_* - u_n\|_E \rightarrow 0$ as $n \rightarrow \infty$ is guaranteed, if there exists a sequence of approximations $(\hat{u}_n)_{n \in \mathbb{N}}$, $\hat{u}_n \in E_n$, satisfying the conditions

$$\|u_* - \hat{u}_n\|_E \rightarrow 0 \text{ as } n \rightarrow \infty, \quad \|A_n^{-1} Q_n A(u_* - \hat{u}_n)\| \rightarrow 0 \text{ as } n \rightarrow \infty.$$

Under two conditions

$$\exists \tau < \infty : \sup_{v_n \in E_n, v_n \neq 0} \frac{\|A v_n\|_F}{\|Q_n A v_n\|_{Z_n^*}} \leq \tau \quad \forall n \in \mathbb{N}, \quad \|A_{n+1}^{-1}\| \text{dist}(f, A E_n) \rightarrow 0 \text{ as } n \rightarrow \infty$$

the self-regularization $\|u_* - u_n\|_E \rightarrow 0$ as $\delta \rightarrow 0$ is guaranteed by choice of the dimension $n(\delta)$ from the discrepancy principle as the first index such that $\|A u_n - f^\delta\|_F \leq b\delta$, $b > \tau + 1$.

In general, estimation of τ may be complicated. We will consider some applications on the collocation method for Volterra integral equations of the first kind.

TIME PERIODIC INVERSE PROBLEM WITH NONLOCAL CONDITION

KRISTINA KAULAKYTĖ

Faculty of Mathematics and Informatics, Vilnius University

Naugarduko str. 24, Vilnius, Lithuania

E-mail: kristina.kaulakyte@mif.vu.lt

In a bounded domain σ we consider the time periodic boundary value problem for the heat equation:

$$\begin{aligned} U_t(x, t) - \nu \Delta U(x, t) &= q(t), \\ U(x, t)|_{\partial\sigma} &= 0, \\ U(x, 0) &= U(x, 2\pi), \\ \int_{\sigma} U(x, t) dx &= F(t), \quad F(0) = F(2\pi), \end{aligned} \tag{1}$$

where U and q are the unknown functions while F is a given function, i.e. for given F we find the right hand side q such that the solution U satisfies the additional nonlocal condition $\int_{\sigma} U(x, t) dx = F(t)$.

Problem (1) can be interpreted as an inverse parabolic problem. Using the concept of a very weak solution introduced in paper [1] for the initial boundary value problem, we prove the existence of a unique weak solution of time periodic problem (1) under the assumption that function F has minimal regularity, i.e function F belongs only to $L^2(0, 2\pi)$ (see [2]).

REFERENCES

- [1] K. Pileckas, R. Čiegis. Existence of nonstationary Poiseuille type solutions under minimal regularity assumptions. *Z. Angew. Math. Phys.*, **71**, 192, 2020. <https://doi.org/10.1007/s00033-020-01422-5>
- [2] K. Kaulakytė, N. Kozulinas and K. Pileckas. Time-periodic Poiseuille-type solution with minimally regular flow rate. *Nonlinear analysis: modelling and control*, **26**, 5, 947–968, 2021. <https://doi.org/10.15388/namc.2021.26.24502>

PERIODIC CUBIC SPLINE HISTOPOLATION

EVELY KIRSIAED, PEETER OJA

University of Tartu

Narva street 18, 51009 Tartu, Estonia

E-mail: evely.kirsiaed@ut.ee, peeter.oja@ut.ee

For a given grid of points $a = x_0 < x_1 < \dots < x_n = b$ and given real numbers z_i , $i = 1, \dots, n$, consider the problem of histopolation, i.e., the problem of finding a function S from some class of functions such that

$$\int_{x_{i-1}}^{x_i} S(x)dx = z_i h_i, \quad i = 1, \dots, n, \quad (1)$$

where $h_i = x_i - x_{i-1}$, $i = 1, \dots, n$.

It is known that cubic spline histopolation problem with classical boundary conditions and arbitrary placement of histopolation knots is uniquely solvable [1]. This is not true for periodic spline histopolation. In case of periodic spline histopolation we look for a solution to the problem from the class of m th degree polynomial splines satisfying boundary conditions

$$S^{(j)}(a) = S^{(j)}(b), \quad j = 0, 1, \dots, m - 1.$$

It is known that for m even, n arbitrary and also for m odd, n odd problem (1) has a unique solution [2]. For m odd, n even it is known that in case of uniform grid there exist values z_i , $i = 1, \dots, n$, such that periodic histopolation problem does not have a solution [2; 3]. In case of non-uniform grid the negative result is proven for $m = 1$, n even and m odd, $n = 2$ (see [2]) leaving other cases open.

We restrict ourselves to the cubic spline case $m = 3$, even number of subintervals $n = 2k$, $k = 2, 3, \dots$ and arbitrary placement of knots; analyse different representations of the histospline and discuss solvability issues.

REFERENCES

- [1] E. Kirsiaed, P. Oja and G. W. Shah. Cubic spline histopolation. *Mathematical Modelling and Analysis*, **22** (4):514–527, 2017.
- [2] P. Oja and G. W. Shah. Periodic polynomial spline histopolation. *Proceedings of the Estonian Academy of Sciences*, **67** (3):246–251, 2018.
- [3] L. Schumaker. *Spline functions: Basic Theory*. Wiley, New York, 1981.

POINTWISE-IN-TIME A-PRIORI AND A-POSTERIORI ERROR CONTROL FOR TIME-FRACTIONAL PARABOLIC EQUATIONS

NATALIA KOPTEVA

University of Limerick

Department of Mathematics and Statistics, University of Limerick, Ireland

E-mail: `natalia.kopteva@ul.ie`

An initial-boundary value problem with a Caputo time derivative of fractional order $\alpha \in (0, 1)$ is considered, solutions of which typically exhibit a singular behaviour at an initial time. For this problem, building on some ideas from [1], we give a simple and general numerical-stability analysis using barrier functions, which yields sharp pointwise-in-time error bounds on quasi-graded temporal meshes with arbitrary degree of grading. This approach is employed in the error analysis of the L1 and Alikhanov L2-1 σ fractional-derivative operators [2], as well as an L2-type discretization of order $3 - \alpha$ in time [3]. This methodology is also generalized for semilinear fractional parabolic equations [4]. In particular, our error bounds accurately predict that milder (compared to the optimal) grading yields optimal convergence rates in positive time. The theoretical findings are illustrated by numerical experiments.

Furthermore, pointwise-in-time a posteriori error bounds will be given in the spatial L_2 and L_∞ norms. Hence, an adaptive mesh construction algorithm is applied for the L1 method, which yields optimal convergence rates $2 - \alpha$ in the presence of solution singularities [5; 6; 7].

REFERENCES

- [1] N. Kopteva. Error analysis of the L1 method on graded and uniform meshes for a fractional-derivative problem in two and three dimensions. *Math. Comp.*, **88** 2135–2155, 2019.
- [2] N. Kopteva and X. Meng. Error analysis for a fractional-derivative parabolic problem on quasi-graded meshes using barrier functions. *SIAM J. Numer. Anal.*, **58** 1217–1238, 2020.
- [3] N. Kopteva. Error analysis of an L2-type method on graded meshes for a fractional-order parabolic problem. *Math. Comp.*, **90** 19–40, 2021.
- [4] N. Kopteva. Error analysis for time-fractional semilinear parabolic equations using upper and lower solutions. *SIAM J. Numer. Anal.*, **58** 2212–2234, 2020.
- [5] N. Kopteva. Pointwise-in-time a posteriori error control for time-fractional parabolic equations. *Appl. Math. Lett.*, **123** 107515, 2022.
- [6] N. Kopteva and M. Stynes. A Posteriori Error Analysis for Variable-Coefficient Multiterm Time-Fractional Subdiffusion Equations. *J. Sci. Comput.*, **92** 73, 2022.
- [7] S. Franz and N. Kopteva. Pointwise-in-time a posteriori error control for higher-order discretizations of time-fractional parabolic equations. *J. Comput. Appl. Math.*, **427** 115122, 2023.

CURVE ESTIMATION WITH MODIFIED COMPLETE SPLINES AND EXPONENTIAL PARAMETERIZATION

RYSZARD KOZERA^{1,2} and MAGDALENA WILKOŁAZKA²

*Warsaw University of Life Sciences - SGGW*¹

*John Paul II Catholic University of Lublin*²

*Institute of Information Technology*¹

*Faculty of Natural and Technical Sciences*²

Nowoursynowska 166, 02-787 Warsaw, Poland¹

Konstantynów 1H, 20-708 Lublin, Poland²

E-mail: ryszard_kozera@sggw.edu.pl, magdalena.wilkolazka@kul.pl

We discuss the issue of fitting data points $\mathcal{Q}_m = \{q_i\}_{i=0}^m$ in arbitrary Euclidean space \mathbb{E}^n (see [1]). It is also assumed, that the knots $\mathcal{T}_m = \{t_i\}_{i=0}^m$ are unknown and as such they need to be replaced by some $\hat{\mathcal{T}}_m = \{\hat{t}_i\}_{i=0}^m$. For \mathcal{Q}_m dense the issue of convergence rate of a given interpolation scheme $\hat{\gamma}$ (based on $\hat{\mathcal{T}}_m$) in approximating γ (with $\gamma(t_i) = q_i$) has been extensively studied (as for classical case with \mathcal{T}_m given - see e.g. [1; 7]). A possible remedy to substitute \mathcal{T}_m with $\hat{\mathcal{T}}_m$ is to apply *an exponential parameterization*:

$$\hat{t}_0 = 0, \quad \hat{t}_{i+1} = \|q_{i+1} - q_i\|^\lambda \quad (1)$$

with $i = 0, \dots, m-1$ and $\lambda \in [0, 1]$ - see [7]. The convergence issue (and its rate) of $\hat{\gamma}$ to γ has been tackled for (1) and $\hat{\gamma}$ forming either *piecewise Lagrange quadratics* or *cubics* or *modified Hermite interpolants* (see [2; 3; 4]). We examine here the asymptotics in $\gamma \approx \hat{\gamma} = \hat{\gamma}^{MC}$ forming a *modified complete spline* based on \mathcal{Q}_m and (1) - see [5]. We establish a *linear convergence rate* (in terms of $\delta_m = \max_{i \in \{0, \dots, m-1\}} \{t_{i+1} - t_i\}$) in estimating γ by $\hat{\gamma}^{MC}$ for $\lambda \in [0, 1)$. The latter supplements the case of $\lambda = 1$ yielding a faster quartic rate. Lastly we report on numerical tests confirming the above asymptotics and *its sharpness*. Finding the knots for *sparse reduced data* (here $m \ll \infty$) can be dealt by solving a relevant optimization task (see [6]).

REFERENCES

- [1] C. de Boor. *A practical guide to spline*. Springer-Verlag, New York Heidelberg Berlin, 1985.
- [2] R. Kozera and L. Noakes. Piecewise-quadratics and exponential parameterization for reduced data. *Applied Mathematics and Computation*, **221**: 620–638, 2013.
- [3] R. Kozera and M. Wilkołazka. Convergence order in trajectory estimation by piecewise-cubics and exponential parameterization. *Mathematical Modelling and Analysis*, **24** (1):72–94, 2019.
- [4] R. Kozera and M. Wilkołazka. A note on modified Hermite interpolation. *Mathematics in Computer Science*, **14**: 223–239, 2020.
- [5] R. Kozera and M. Wilkołazka. A modified complete spline interpolation and exponential parameterization. In: *Proc. of the 14th Intern. Conference CISIM-2015, Warsaw, Poland, 2015*, Federation for Information Processing, K. Saeed and W. Homenda (Eds.), Cham Springer, Vol. 9339, 2015, 98 – 110.
- [6] R. Kozera, L. Noakes and A. Wiliński. Generic case of Leap-Frog algorithm for optimal knots selection in fitting reduced data. In: *Proc. of the 21st Intern. Conference ICCS-2021, Kraków, Poland, 2021*, Computational Science, M. Paszyński, D. Kranzlmüller and V.V. Krzhizhanovskaya (Eds.), Cham Springer, Vol. 12745 Part IV, 2021, 337 – 350.
- [7] B.I. Kvasov. *Methods of shape-preserving spline approximation*. World Scientific, Singapore, 2000.

BLOOD FLOW MODELLING IN HUMAN HEART USING FSI

NIKOLAJUS KOZULINAS

Vilnius University

Naugarduko str. 24, 03225 Vilnius, Lithuania

E-mail: `nikolajus.kozulinas@mif.vu.lt`

The goal of this talk is to present the FSI (fluid-structure interaction) CFD (computational fluid dynamics) simulations of the blood flow in the human heart left atrial appendage for a patient-specific geometry. These simulations are important for the medical doctors decision making for the patients with atrial fibrillation. In order to compute blood flow velocity we use Navier-Stokes equations coupled with Shell mechanics Uflyand-Mindlin model. The FSI CFD simulations in the heart is a challenging problem: the existing softwares are not too robust for real life Reynolds numbers and often do not converge to the solution of the Navier-Stokes equations for the blood coupled with the elasticity equations of the wall. That is why we first provide the CFD computations with the rigid wall when the codes are more stable. Using this solution as the reference velocity and pressure, then we provide the FSI computations which become much more robust (see [1]).

REFERENCES

- [1] A. Aidietis, S. Aidietienė, O. Ardatov, S. Borodinas, R. Katkus, K. Kaulakytė, N. Kozulinas, G. Panasenko, K. Pileckas. Efficient blood velocity computation in human heart left atrial appendage. *IN PREPARATION*, 2024.

ABOUT THE NUMERICAL SOLUTION OF SINGULAR FRACTIONAL INTEGRO-DIFFERENTIAL EQUATIONS

KAIDO LÄTT

University of Tartu

Narva street 18, 51009 Tartu, Estonia

E-mail: `kaido.latt@ut.ee`

Due to various new applications in physics, chemistry, and other fields of science, the interest in fractional derivatives and equations containing them has increased significantly over last century (see for example [1; 3]). In [2] a class of singular fractional integro-differential equations was investigated.

We now consider singular fractional integro-differential equations of the form

$$(M^\beta D_0^\beta u)(t) = a_1(M^{\beta_1} D_0^{\beta_1} u)(t) + a_0(V_\varphi u)(t) + f(t), \quad 0 < t \leq T. \quad (1)$$

By $C^m[0, T]$ ($m \in \mathbb{N}_0$) we denote the space of m times continuously differentiable functions u on the closed interval $[0, T]$; $C^0[0, T] = C[0, T]$, and by $L^1(0, 1)$ we denote the Banach space consisting of real or complex valued functions φ defined on $(0, 1)$ such that $\|\varphi\|_{L^1(0,1)} = \int_0^1 |\varphi(x)| dx < \infty$. In equation (1), the operator M^β ($\beta \in \mathbb{R}$) is defined as $(M^\beta u)(t) = t^\beta u(t)$ ($0 < t \leq T$) for $u \in C[0, T]$,

$$(V_\varphi u)(t) = \int_0^t \frac{1}{t} \varphi\left(\frac{s}{t}\right) u(s) ds = \int_0^1 \varphi(x) u(tx) dx, \quad 0 \leq t \leq T, \quad u \in C[0, T],$$

i.e. $V_\varphi : C[0, T] \rightarrow C[0, T]$ is a cordial Volterra integral operator with core $\varphi \in L^1(0, 1)$, and

$$\beta, \beta_1, a_0, a_1 \in \mathbb{R}, \quad q < \beta < q + 1, \quad \beta > \beta_1 \geq 0, \quad f \in C^q[0, T].$$

The fractional differential operator D_0^β (of the order $\beta \in [0, \infty)$) in equation (1), is defined as the inverse of the Riemann-Liouville integral operator $J^\beta : C[0, T] \rightarrow C[0, T]$ on the space $J^\beta C[0, T]$, i.e. $D_0^\beta v = (J^\beta)^{-1}v$, where v belongs to the range $J^\beta C[0, T]$ of J^β , $\beta \geq 0$.

We study the unique solvability of equations of the form (1) and discuss the numerical solution of such equations.

This is a joint work with Arvet Pedas.

REFERENCES

- [1] K. Diethelm. *The Analysis of Fractional Differential Equations*. Springer-Verlag, Berlin, 2010.
- [2] K. Lätt and A. Pedas. Numerical schemes for a class of singular fractional integro-differential equations. *Applied Numerical Mathematics*, **200** 331–343, 2024.
- [3] I. Podlubny. *Fractional Differential Equations*. Academic Press, San Diego, 1999.

APPROXIMATION BY BEURLING'S ZETA-FUNCTIONS

ANTANAS LAURINČIKAS

Vilnius University

Naugarduko street 24, LT-03225 Vilnius, Lithuania

E-mail: `antanas.laurincikas@mif.vu.lt`

The system \mathcal{P} of real numbers $1 < p_1 \leq p_2 \leq \dots \leq p_n \rightarrow \infty$ as $n \rightarrow \infty$, is called generalized prime numbers. Using the system \mathcal{P} , the system $\mathcal{N}_{\mathcal{P}}$ of generalized integers

$$m = p_1^{\alpha_1} \cdots p_r^{\alpha_r}, \quad r \in \mathbb{N}, \quad \alpha_j \in \mathbb{N}_0 = \mathbb{N} \cup \{0\}, \quad j = 1, \dots, r,$$

is constructed. Generalized prime numbers were introduced by Beurling [1], he also obtained the first results on the asymptotic behaviour as $x \rightarrow \infty$ for

$$\pi_{\mathcal{P}}(x) = \sum_{p \leq x, p \in \mathcal{P}} 1 \quad \text{and} \quad \mathcal{N}_{\mathcal{P}}(x) = \sum_{m \leq x, m \in \mathcal{N}_{\mathcal{P}}} 1,$$

and relations among them. To study $\pi_{\mathcal{P}}(x)$, Beurling introduced zeta-functions $\zeta_{\mathcal{P}}(s)$ defined in some half-planes by

$$\zeta_{\mathcal{P}}(s) = \prod_{p \in \mathcal{P}} \left(1 - \frac{1}{p^s}\right)^{-1} \quad \text{or} \quad \zeta_{\mathcal{P}}(s) = \sum_{m \in \mathcal{N}_{\mathcal{P}}} \frac{1}{m^s}.$$

Now, the functions $\zeta_{\mathcal{P}}(s)$ are called Beurling zeta-functions. Various authors developed analytic theory for the functions $\zeta_{\mathcal{P}}(s)$. Note that each case of \mathcal{P} requires a separate studying.

In report, we consider the approximation problem of analytic functions by shifts $\zeta_{\mathcal{P}}(s + i\tau)$ for some classes of systems \mathcal{P} . We assume the estimation

$$\mathcal{N}_{\mathcal{P}}(x) = ax + O(x^{\delta}), \quad 0 \leq \delta < 1, \quad a > 0,$$

and suppose that there exists $\hat{\sigma} < 1$, such that

$$\hat{\sigma} = \inf \left\{ \sigma : \int_0^T |\zeta_{\mathcal{P}}(\sigma + it)|^2 dt \ll_{\sigma} T, \quad \sigma > \delta \right\}.$$

Under the above hypotheses, we obtain [2] that there exists a set of analytic functions, approximated by shifts $\zeta_{\mathcal{P}}(s + i\tau)$. Identification of that set requires new restrictions for the system \mathcal{P} .

REFERENCES

- [1] A. Beurling. Analyse de la loi asymptotique de la distribution des nombres premiers généralisés. I. *Acta Math.*, **68** :225–291, 1937.
- [2] A. Laurinčikas. On value distribution of certain Beurling zeta-functions. *Mathematics*, **12** (3):article no. 459, 2024.

ON THE NUMERICAL SOLUTION OF THE NEUMANN PROBLEM FOR LAPLACE'S EQUATION IN PLANAR DOMAINS WITH CORNERS

PETER JUNGHANNS, CONCETTA LAURITA

Department of Mathematics, Chemnitz University of Technology

Reichenhainer Str. 41, 09126 Chemnitz, Germany

Department of Mathematics, Computer Science and Economics, University of Basilicata

Viale dell'Ateneo Lucano n. 10, 85100 Potenza, Italy

E-mail: `concetta.laurita@unibas.it`

A new boundary integral equation (BIE) method for the numerical solution of the exterior Neumann problem for the Laplace equation in planar domains with corners is proposed. Using the single layer representation of the potential, the differential problem is reformulated in terms of a BIE whose solution has singularities at the corners.

A "modified" Nyström type method based on a Gaussian type quadrature formula is proposed for its approximation.

Convergence and stability results for the proposed method are proved in proper weighted spaces of continuous functions. Moreover, the use of a smoothing transformation allows to increase the regularity of the solution and, consequently, the order of convergence of the method.

The efficiency of the proposed method is shown by illustrating some numerical tests.

ON CENTRAL PART INTERPOLATION APPROXIMATIONS FOR SYSTEMS OF FRACTIONAL DIFFERENTIAL EQUATIONS

MARGUS LILLEMÄE

University of Tartu

Narva street 18, 51009 Tartu, Estonia

E-mail: `margus.lillemae@ut.ee`

We consider an initial value problem for a system of fractional differential equations of the form

$$\begin{aligned} (D_{Cap}^\alpha y_i)(t) + \sum_{j=1}^n a_{ij}(t)y_j(t) &= f_i(t), \quad 0 \leq t \leq 1, \quad i = 1, \dots, n, \quad n \in \mathbb{N} := \{1, 2, \dots\}, \quad (1) \\ y_i(0) &= y_{0i}, \quad y_{0i} \in \mathbb{R} := (-\infty, \infty), \quad i = 1, \dots, n, \quad (2) \end{aligned}$$

where $0 < \alpha < 1$, and the functions a_{ij}, f_i are continuous: $a_{ij}, f_i \in C[0, 1]$ for $i, j = 1, \dots, n$. Here $D_{Cap}^\alpha y$ is an α -order Caputo fractional derivative of a function $y = y(t)$. Then it turns out that problem (1)-(2) possesses a unique solution y_1, \dots, y_n such that $y_i \in C[0, 1]$, $D_{Cap}^\alpha y_i \in C[0, 1]$, $i = 1, \dots, n$. However, we cannot generally expect (see [1]) that y_1, \dots, y_n belong to $C^q[0, 1]$ for $a_{ij}, f_i \in C^q[0, 1]$, $i, j = 1, \dots, n$, $q \in \mathbb{N}$. Instead, we can show that, for $i, j = 1, \dots, n$, if $a_{ij}, f_i \in C^{q,\mu}(0, 1]$, $q \in \mathbb{N}$, $\mu \in (0, 1)$, then y_i and its derivatives $D_{Cap}^\alpha y_i$ belong to $C^{q,\nu}(0, 1]$, where $\nu = \max\{1 - \alpha, \mu\}$. Here, by $C^{q,\mu}(0, 1]$ ($q \in \mathbb{N}$, $0 < \mu < 1$) we denote the set of functions $y \in C[0, 1] \cap C^q(0, 1]$ such that

$$|y^{(k)}(t)| \leq c t^{1-\mu-k}, \quad 0 < t \leq 1, \quad k = 1, \dots, q,$$

where c is a positive constant independent of t .

We propose a high-order method for solving (1)-(2) based on improving the boundary behavior of the exact solution with the help of a change of variables, and on central part interpolation by polynomial splines on the uniform grid. The central part interpolation approach was used for solving Fredholm integral equations of the second kind and it has shown accuracy and numerical stability advantages compared to standard piecewise polynomial collocation methods, including collocation at Chebyshev points [2]. We apply this approach for solving (1)-(2) and derive global error estimates for the approximate solution analogous to [3].

This work is my joint collaboration with Arvet Pedas and Mikk Vikerpuur.

REFERENCES

- [1] A. Pedas, M. Vikerpuur. Spline collocation for multi-term fractional integro-differential equations with weakly singular kernels. *Fractal Fract.*, **5** 90, 2021.
- [2] K. Orav-Puurand, A. Pedas, G. Vainikko. Central part interpolation schemes for integral equations with singularities. *J. Integral Equ. Appl.*, **29** 401–440, 2017.
- [3] M. Lillemäe, A. Pedas, M. Vikerpuur. Central part interpolation schemes for fractional differential equations. *Applied Numerical Mathematics*, **200** 318–330, 2024.

MATHEMATICAL MODELING OF NEURAL FIELDS WITH DIFFUSION

DANIELE AVITABILE (1), NIKOLAI CHEMETOV (2), AND PEDRO LIMA (3)

*Amsterdam Centre for Dynamics and Computations, Vrije Universiteit Amsterdam(1);
DCM-FFCLRP, University of S. Paulo(2); CEMAT, Instituto Superior Tecnico, University of
Lisbon (3)*

Amsterdam, Holland (1); Ribeirão Preto, Brasil (2); Lisboa, Portugal (3)

E-mail: `plima@math.tecnico.ulisboa.pt`

We are concerned with the analytical and numerical solution of the following integro-differential equation

$$\begin{aligned} \partial_t v(x, \xi, t) &= \nu \partial_{\xi\xi}^2 v(x, \xi, t) - \gamma v(x, \xi, t) \\ &+ \int_{\Omega} W(x, x', \xi, \xi') S(v(x', \xi', t)) dx' d\xi' + G(x, \xi, t), \end{aligned} \quad (1)$$

for $(x, \xi, t) \in \Omega_T = \Omega \times [0, T]$, where $\Omega = \mathbb{T}^n \times U$, $U = (0, L)$ and \mathbb{T}^n is the n -dimensional torus.

We search for a solution v of (1), being a periodic function on x , satisfying Neumann boundary conditions with respect to ξ and an initial condition.

Our analysis, which is described with detail in [2], relies on perturbing weak solutions to the diffusion-less problem (with $\nu = 0$), that is, a standard neural field, for which weak problems have not been studied to date. We find rigorous asymptotic estimates for the problem with and without diffusion, and prove that the solutions of the two models stay close, in a suitable norm, on finite time intervals. Using a computational method that is described in [1], we provide numerical evidence of our perturbative results.

REFERENCES

- [1] D. Avitabile, S. Coombes, and P. Lima. Numerical investigation of a neural field model including dendritic processing. *Journal of Computational Dynamics*, **7** 270-290, 2020.
- [2] D. Avitabile, N. Chemetov and P. Lima. Well-posedness and regularity of solutions to neural field problems with dendritic processing. *submitted for publication*, .

SOME GENERALIZED LIMIT THEOREMS FOR THE EPSTEIN ZETA-FUNCTION

RENATA MACAITIENĖ

Institute of Regional Development, Šiauliai Academy, Vilnius University

Vytauto str. 84, LT-76352 Šiauliai, Lithuania

Faculty of Business and Technologies, Šiauliai State Higher Education Institution

Aušros av. 40, LT-76241 Šiauliai, Lithuania

E-mail: `renata.macaitiene@sa.vu.lt`

Let Q be a positive definite quadratic $n \times n$ matrix and $Q[\underline{x}] = \underline{x}^T Q \underline{x}$ for $\underline{x} \in \mathbb{Z}^n$. The Epstein zeta-function $\zeta(s; Q)$, $s = \sigma + it$, is defined, for $\sigma > \frac{n}{2}$, by the series $\zeta(s; Q) = \sum_{\underline{x} \in \mathbb{Z}^n \setminus \{0\}} (Q[\underline{x}])^{-s}$, and can be continued analytically to the whole complex plane, except for a simple pole at the point $s = \frac{n}{2}$ with residue $\frac{\pi^{n/2}}{\Gamma(n/2)\sqrt{\det Q}}$. The function $\zeta(s; Q)$ was introduced by P. Epstein in [1], its value-distribution was investigated by various authors; for example, an extensive survey of the results for the function $\zeta(s; Q)$ is given in [2].

In the talk, some generalized continuous and discrete results of joint works with A. Laurinčikas on the value-distribution for $\zeta(s; Q)$ with even $n \geq 4$ and integers $Q[\underline{x}]$ will be presented. More precisely, we will show that, for a special differentiable function $\varphi(t)$ with a monotonic derivative,

$$\frac{1}{T} \text{meas} \{t \in [0, T] : \zeta(\sigma + i\varphi(t); Q) \in A\}, \quad A \in \mathcal{B}(\mathbb{C}),$$

converges weakly to an explicitly given probability measure on $(\mathbb{C}, \mathcal{B}(\mathbb{C}))$ as $T \rightarrow \infty$ [3]. Here $\text{meas}A$ denotes the Lebesgue measure of a measurable set $A \subset \mathbb{R}$, and $\mathcal{B}(\mathbb{C})$ – the Borel σ -field of the space \mathbb{C} . Moreover, we will discuss that, for a certain increasing differentiable function $\varphi(t)$ with a continuous monotonic bounded derivative and with an additional condition for the sequence $\{\varphi(k)\}$, on $(\mathbb{C}, \mathcal{B}(\mathbb{C}))$, there exists an explicitly described probability measure $P_{Q, \sigma}$ such that

$$\frac{1}{N} \# \{N \leq k \leq 2N : \zeta(\sigma + i\varphi(k); Q) \in A\}, \quad A \in \mathcal{B}(\mathbb{C}),$$

converges weakly to $P_{Q, \sigma}$ as $N \rightarrow \infty$ [4]. Also, a few examples of the function $\varphi(t)$ will be given.

REFERENCES

- [1] P. Epstein. Zur Theorie allgemeiner Zetafunktionen. *Math. Ann.*, **56**: 615–644, 1903.
- [2] T. Nakamura, L. Pańkowski. On zeros and c -values of Epstein zeta-functions. *Šiauliai Math. Semin.*, **8** (16): 181–195, 2013.
- [3] A. Laurinčikas, R. Macaitienė. A generalized Bohr–Jessen type theorem for the Epstein zeta-function. *Mathematics*, **10** (12): 2042, 1–11, 2022.
- [4] A. Laurinčikas, R. Macaitienė. A generalized discrete Bohr–Jessen type theorem for the Epstein zeta-function. *Mathematics*, **11** (4): 799, 1–13, 2023.

ON MATHEMATICAL MODELLING AND SIMULATION OF IMMUNE SYSTEM'S RESPONSE IN TUMOR TREATMENT

MAKSIMS MARINAKI^{1,2}, MANFREDS ŠNEPS-ŠNEPPE³

Institute of Mathematics and Computer Science¹, University of Latvia², Ventspils University of Applied Sciences³

Raiņa bulvāris 29, Rīga, LV-1459, Latvija¹

Jelgavas iela 3, Rīga, LV-1004, Latvija²

Inženieru iela 101, Ventspils, LV-3601, Latvija³

E-mail: maksims.marinaki@lu.lv

E-mail: manfreds.sneps@venta.lv

We address the intricacies surrounding the efficacy of oncolytic viruses and the infiltration of healthy cells to combat malignant cells. Similar models and treatment of experimental data for the process also explored in [1].

We consider PDE systems (1) with special right hand sides modelling the different amounts of injections:

$$\begin{cases} \frac{\partial C}{\partial t} - D\Delta C = \sum \gamma\delta(x - x_p) + f(x), & x \in \Omega, \\ D\frac{\partial C}{\partial n} + \beta C = 0, & x \in \partial\Omega, \end{cases} \quad (1)$$

where $\delta(x)$ is the dirac delta function, C is the concentration, D is the diffusivity, γ is the injection rate and x_p is the injection position.

These are discretized by finite element techniques.

Consideration of simpler models for similar problems in the field such as ODE systems has been explored in [2] and main features are discussed here as well. We present the graphical results of the obtained fields and discuss main aspects of several underlying modelling stages. Moreover we discuss the presence of experimental data and the identification of parameters for these models.

REFERENCES

- [1] Chen, J., Weihs, D., Vermolen, F.J. A Cellular Automata Model of Oncolytic Virotherapy in Pancreatic Cancer. *Bull Math Biol*, **82**, 2020.
- [2] Li, X., Xu, J.X. A Mathematical Model of Immune Response to Tumor Invasion Incorporated with Danger Model. *Journal of Biological Systems*, **23** (3), 505 – 526, 2015.

ON GENERALIZED SHIFTS OF THE LERCH ZETA-FUNCTION

TOMA MIKALAUŠKAITĖ

Vilnius University

Naugarduko street 24, LT-03225 Vilnius, Lithuania

E-mail: toma.mikalauskaite@mif.stud.vu.lt

Let $s = \sigma + it$ be a complex variable, and $0 < \lambda \leq 1$, $0 < \alpha \leq 1$ two parameters. The Lerch zeta-function $L(\lambda, \alpha, s)$ is defined, for $\sigma > 1$, by the Dirichlet series

$$L(\lambda, \alpha, s) = \sum_{m=0}^{\infty} \frac{e^{2\pi i \lambda m}}{(m + \alpha)^s},$$

and has a meromorphic continuation to the whole complex plane with a simple pole at the point $s = 1$ when $\lambda = 1$.

In the report, we consider the approximation of analytic functions by generalized shifts $L(\lambda, \alpha, s + i\varphi(\tau))$. The real-valued function $\varphi(\tau)$ is defined for $\tau \geq \tau_0 > 0$, is increasing to $+\infty$ and has a monotonic derivative such that

$$\varphi(2\tau) (\min(\varphi'(\tau), \varphi'(2\tau))) = O(\tau), \quad \tau \rightarrow \infty.$$

We present the following results.

THEOREM 1. *Suppose that the set $\{\log(m + \alpha) : m \in \mathbb{N}_0\}$ is linearly independent over the field of rational numbers, K is a compact subset of the strip $D = \{s \in \mathbb{C} : 1/2 < \sigma < 1\}$ with connected complement, and $f(s)$ is a continuous on K function, and analytic in the interior of K . Then, for every $\varepsilon > 0$,*

$$\liminf_{T \rightarrow \infty} \frac{1}{T} \text{meas} \left\{ \tau \in [0, T] : \sup_{s \in K} |L(\lambda, \alpha, s + i\varphi(\tau)) - f(s)| < \varepsilon \right\} > 0. \quad (1)$$

Moreover, “ \liminf ” can be replaced by “ \lim ” for all but at most countably many $\varepsilon > 0$.

THEOREM 2. *Suppose that the parameters λ and α are arbitrary. Then there exists a non-empty closed set $F_{\lambda, \alpha, \varphi} \subset D$ such that, for every compact set $K \subset D$, $f(s) \in F_{\lambda, \alpha, \varphi}$ and $\varepsilon > 0$, inequality (1) is valid, and the same assertion as in Theorem 1 with “ \lim ” is true..*

Theorems 1 and 2 generalize the results of [1] and [2] obtained for $\varphi(\tau) = \tau$.

REFERENCES

- [1] A. Laurinćikas. The universality of the Lerch zeta-function. *Lith. Math. J.*, **37** (3):275–280, 1997.
- [2] A. Laurinćikas. “Almost” universality of the Lerch zeta-function. *Math. Commun.*, **24** (1):107–118, 2019.

STABILITY ANALYSIS OF STATIONARY MOTION OF THE 3D SWINGING ATWOOD'S MACHINE

ALEXANDER PROKOPENYA

Institute of Information Technology, Warsaw University of Life Sciences – SGGW

Nowoursynowska street 159, 02-776 Warsaw, Poland

E-mail: alexander_prokopenya@sggw.edu.pl

The swinging Atwood machine under consideration consists of two masses m_1 , m_2 attached to opposite ends of a massless inextensible thread wound round two massless frictionless pulleys of negligible radius (see [1]). The mass m_1 is allowed to swing in two dimensions and its behaves like a spherical pendulum of variable length while the mass m_2 is constrained to move only along a vertical. Such a system has three degrees of freedom and its equations of motion may be written in the form

$$\begin{aligned} r\ddot{\theta} &= -g \sin \theta - 2\dot{r}\dot{\theta} + \frac{p_\varphi^2 \cos \theta}{r^3 \sin^3 \theta}, & r^2 \sin^2 \theta \dot{\varphi} &= p_\varphi = \text{const}, \\ (2 + \varepsilon)\dot{r} &= r\dot{\theta}^2 - g(1 + \varepsilon - \cos \theta) + \frac{p_\varphi^2}{r^3 \sin^2 \theta}. \end{aligned} \quad (1)$$

Here the variables r, θ, φ describe geometrical configuration of the system, g is a gravity constant, $\varepsilon = (m_2 - m_1)/m_1$, and p_φ is an integral of motion determined from the initial conditions.

In the case of $p_\varphi = 0$ the mass m_1 oscillates on a vertical plane and we obtain the swinging Atwood machine which may demonstrate a periodic motion (see [2; 3]).

One can easily check that there exists an exact particular solution to equations (1) of the form

$$\varphi(t) = \sqrt{\frac{g(1 + \varepsilon)}{r_0}} t + \varphi_0, \quad r(t) = r_0, \quad \theta(t) = \theta_0 = \arccos(1/(1 + \varepsilon)). \quad (2)$$

Solution (2) describes a uniform motion of the mass m_1 in a horizontal plane on a circular orbit of radius $r_0 \sin \theta_0$. Simulation of the system motion shows that small variation of the initial conditions results only in small perturbation of the mass m_1 orbit. Analyzing the Hamiltonian function of the system and applying the stability theory, we have proved that solution (2) is stable with respect to the variables $r, \dot{r}, \theta, \dot{\theta}, \varphi$.

REFERENCES

- [1] A.N. Prokopenya. Motion of a swinging Atwood's machine: simulation and analysis with Mathematica. *Mathematics in Computer Science*, **11** (3-4):417–425, 2017.
- [2] A.N. Prokopenya. Construction of a periodic solution to the equations of motion of generalized Atwood's machine using computer algebra. *Programming and Computer Software*, **46** (2):120–125, 2020.
- [3] A. Prokopenya. Stability analysis of periodic motion of the swinging Atwood machine. In: *Proc. of the 24th Intern. Workshop CASC 2022, Gebze, Turkey, 2022*, Computer Algebra in Scientific Computing, F. Boulrier, M. England, T.M. Sadykov, E.V. Vorozhtsov (Eds.), LNCS 13366. Springer, Cham, 2022, 288 – 299.

ON THE CONVERGENCE OF THE DIFFERENCE SCHEME FOR NONLINEAR PARABOLIC EQUATION WITH INTEGRAL BOUNDARY CONDITION

K. PUPALAIĖ^{1,2}, M. SAPAGOVAS³

¹*Department of Applied Mathematics, Kaunas University of Technology*

Studentų str. 50, LT-51368, Kaunas, Lithuania

²*Kaunas University of Applied Engineering Sciences*

Tvirtovės al. 35, LT-50155, Kaunas, Lithuania

E-mail: kristina.pupalaige@ktu.lt

³*Institute of Data Science and Digital Technologies, Vilnius University*

Akademijos str. 4, LT-08412, Vilnius, Lithuania

E-mail: mifodijus.sapagovas@mif.vu.lt

We analyze the two dimensional nonlinear parabolic equation

$$\frac{\partial u}{\partial t} = \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} - f(x, y, u), \quad (x, y) \in \Omega = \{0 < x < 1, 0 < y < 1\}, \quad t \in (0, T],$$

with given initial condition and boundary conditions

$$\begin{aligned} u(x, y, 0) &= \phi(x, y), \quad (x, y) \in \Omega, \\ u(x, 0, t) &= \mu_1(x, t), \quad u(x, 1, t) = \mu_2(x, t), \quad u(0, y, t) = \mu_3(y, t) \end{aligned}$$

and with nonlocal integral condition on one boundary of rectangular domain

$$u(1, y, t) = \gamma \int_0^1 \int_0^1 u(x, y, t) dx dy + \mu_4(y, t).$$

We write down the finite difference method and estimate the error of the solution based on the properties of M-matrices, and then prove the convergence of the differential scheme. The main result of the investigation is that the majorant function is constructed and establishing the stability and convergence of the differential scheme.

REFERENCES

- [1] K. Bingelė, and A. Bankauskienė, and A. Štikonas. Spectrum curves for a discrete Sturm-Liouville problem with one integral boundary condition. *Nonlin. Anal. Model. Control.*, **24** 755–744, 2019.
- [2] M. Sapagovas, O. Štikonienė, K. Jakubėlienė, R. Čiupaila. Finite difference method for boundary value problem for nonlinear elliptic equation with nonlocal conditions. *Bound. Value Probl.*, **2019** (94):1–16, 2019.
- [3] K. Pupalaigė, and M. Sapagovas", and R. Čiupaila. Nonlinear elliptic equation with nonlocal integral boundary condition depending on two parameters. *Math. Model. Anal.*, **27** (4), 610–628, 2022.
- [4] M. Sapagovas, and J. Novickij. On stability in the maximum norm of difference scheme for nonlinear parabolic equation with nonlocal condition. *Nonlin Anal. Model Contr.*, **28** 365-376, 2023.

SOLITARY SOLUTIONS TO THE HEPATITIS C MODEL WITH THE PROLIFERATION OF INFECTED AND UNINFECTED HEPATOCYTES

MINVYDAS RAGULSKIS

Department of mathematical modelling, Kaunas University of Technology

Studentu 50-146, Kaunas LT-51368, Lithuania

E-mail: minvydas.ragulskis@ktu.lt

It is shown in [1] that the model of the Hepatitis C model with the proliferation of infected and uninfected hepatocytes [2] can be expressed in the form of two Riccati equations coupled with the multiplicative and diffusive terms:

$$\begin{cases} \frac{dx}{dt} = a_0 + a_1x + a_2x^2 + a_3xy + a_4y, \\ \frac{dy}{dt} = b_0 + b_1y + b_2y^2 + b_3xy + b_4x, \end{cases} \quad (1)$$

where $a_k, b_k \in \mathbb{R}; k = 0, 1, \dots, 4$. The standard form of a solitary solution reads:

$$x(t) = \sigma \frac{\prod_{k=1}^n (\exp(\eta(t-t_0)) - x_k)}{\prod_{k=1}^n (\exp(\eta(t-t_0)) - t_k)}, \quad (2)$$

where $n \in \mathbb{N}$ is the order of the solitary solution; $\sigma, \eta, t_0, x_k, t_k \in \mathbb{R}$.

It is a common feature of nonlinear dynamical systems that a solitary solution form a separatrix in the space of system parameters and initial conditions [3]. Therefore, the existence of solitary solutions may help to understand the global dynamics of the system. The necessary condition for the existence of the order-1 solitary solutions to Eq.(1) is derived in [1]:

$$a_3 = b_2; b_3 = a_2. \quad (3)$$

However, the limit transition from Eq.(1) to the system of Riccati equations coupled with the diffusive terms yields a system of degenerate Riccati equations [4]. Moreover, the fact that the Riccati system coupled with the diffusive terms admits non-deformed order-1 solitary solutions, serves as a proof for the structural instability of the hepatitis C model with the proliferation of infected and uninfected hepatocytes.

REFERENCES

- [1] T. Telksnys, Z. Navickas, M.A.F. Sanjuán, R. Marcinkevicius and M. Ragulskis. Kink solitary solutions to a hepatitis C evolution model. *Discrete And Continuous Dynamical Systems-B*, **23** 4427–4447, 2020.
- [2] T.C. Reluga, H. Dahari and A.S. Perelson. Analysis of hepatitis C virus infection models with hepatocyte homeostasis. *SIAM Journal of Applied Mathematics*, **69** (4):999–1023, 2009.
- [3] T. Telksnys, Z. Navickas, I. Timofejeva, R. Marcinkevicius and M. Ragulskis. Symmetry breaking in solitary solutions to the Hodgkin–Huxley model. *Nonlinear Dynamics*, **97** (1):571–582, 2019.
- [4] Z.Navickas, R.Marcinkevicius, I.Telksniene, T.Telksnys and M.Ragulskis. Structural stability of the Hepatitis C model with the proliferation of infected and uninfected hepatocytes. *Mathematical and Computer Modelling of Dynamical Systems*, **30** (1):51–72, 2024.

NEW HEURISTIC PARAMETER CHOICE FOR THE CLASS OF REGULARIZATION METHODS

TOOMAS RAUS

Institute of Mathematics and Statistics, University of Tartu

Narva mnt 18, Tartu, 51009, Estonia

E-mail: toomas.raus@ut.ee

We consider an operator equation

$$Au = f_*, \quad f_* \in R(A) \neq \overline{R(A)},$$

where $A \in L(H, F)$ is the linear continuous operator between real Hilbert spaces H and F . Instead of the exact right-hand side f_* we have only an approximation $f \in F$. To get the regularized solution we consider the following class of regularization methods

$$u_r = (I - A^* A g_r(A^* A)) u_0 + g_r(A^* A) A^* f,$$

where u_0 is the initial approximation, r is the regularization parameter and the generating function $g_r(\lambda)$ satisfies for $r \geq 0$ the conditions

$$\sup_{0 \leq \lambda \leq \|A^* A\|} |g_r(\lambda)| \leq \gamma r, \quad \sup_{0 \leq \lambda \leq \|A^* A\|} \lambda^p |1 - \lambda g_r(\lambda)| \leq \gamma_p r^{-p}, \quad 0 \leq p \leq p_0.$$

Well-known heuristic rule for choosing regularization parameter is the quasioptimality principle, where the parameter is chosen as the global minimum point of the function

$$\psi_Q(r) = r \|A^* B_r^2 (A u_r - f)\|, \quad B_r = (I - A A^* g_r(A A^*))^{1/2p_0},$$

on the set of parameters $\Omega = \{r_j : r_j = q r_{j-1}, j = 1, 2, \dots, M, q > 1\}$. Unfortunately, this rule is unstable in this sense that it often fails in case of heat-type problems.

To get stable parameter choice rule we introduce modified quasioptimality criterion function $\psi_{MQ}(r)$ in the form:

$$\psi_{MQ}(r_0) = \psi_Q(r_0),$$

$$\psi_{MQ}(r_j) = \max\{\psi_Q(r_j), (d_{MD}(r_j)/d_{MD}(r_{j-1}))^{2/(2+1/p_0)} \psi_{MQ}(r_{j-1})\}, j > 0,$$

where the function $d_{MD}(r) = \|B_r(A u_r - f)\|$. Choosing global minimum point of the function $\psi_{MQ}(r)$ as regularization parameter gives us stable heuristic rule, but to get more accuracy we present the following algorithm. Let r_{MQ} be the global minimum point of the function $\psi_{MQ}(r)$ on the set of local minimum points of the function $\psi_Q(r)$.

Heuristic rule. For the regularization parameter choose the parameter

$$r_H = \min\{r_Q, \max\{r_{MQ}, r_{HR}\}\},$$

where r_Q and r_{HR} are the global minimum points of the functions $\psi_Q(r)$ and $\psi_{HR}(r) = r^{1/2} d_{MD}(r)$, respectively.

A POLYNOMIAL COLLOCATION METHOD FOR THE Lighthill's PROBLEM

MAGDA REBELO

Center for Mathematics and Applications (NOVA Math) and Department of Mathematics, NOVA FCT.

Quinta da Torre, 2829-516, Caparica, Portugal

E-mail: msjr@fct.unl.pt

Lighthill [1] derived a nonlinear integral equation which describes the temperature distribution of the surface of a projectile moving through a laminar layer at high Mach numbers, which is given by

$$2t^{1/3}u^4(t) + \Gamma(2/3) \left(J^{2/3}u' \right) (t) = 0, t > 0, \quad (1)$$

$$u(0) = 1, \quad (2)$$

where J^α , $0 < \alpha < 1$, is the Riemann-Liouville operator defined by

$$(J^\alpha v)(t) = \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} v(s) ds, \quad t \geq 0.$$

In [2] was proved that (1)-(2) possesses a unique continuous solution, u^* , such that

$$0 < u^*(t) < 1, \quad t > 0; \quad u^*(t) \longrightarrow 0 \text{ as } t \rightarrow +\infty,$$

with the following unimprovable, for large t , estimate

$$u^*(t) \leq \frac{\sin(\pi/3)}{\sqrt{3}} t^{-1/6} \sim 0.841t^{-1/6}.$$

Then, the convergence $u^*(t) \longrightarrow 0$ as $t \rightarrow +\infty$ is slow.

This work is concerned with the numerical solution of (1). Indeed we consider a numerical method based on polynomial collocation method, with several choices for the mesh, and compare the numerical results obtained.

Some considerations about the convergence of the proposed method are made.

REFERENCES

- [1] Lighthill, M. J. . Contributions to the theory of heat transfer through a laminar boundary layer . *Proceedings of the Royal Society of London. Series A. Mathematical and Physical Sciences* , **202** (1070): 359–377, 1950.
- [2] Vainikko, G. . Positive Solution of Lighthill-Type Equations . *Zeitschrift für Analysis und ihre Anwendungen*, **37** (4): 475–494, 2018 .

DISCRETE VERSION OF THE MISHOU THEOREM FOR ABSOLUTELY CONVERGENT DIRICHLET SERIES RELATED TO PERIODIC ZETA-FUNCTIONS

AIDAS BALČIŪNAS, AUDRONE RIMKEVIČIENĖ

Institute of Mathematics, Faculty of Mathematics and Informatics, Vilnius University
Faculty of Business and Technologies, Šiauliai State University of Applied Sciences

Naugarduko str. 24, LT-03225 Vilnius, Lithuania

Aušros av. 40, LT-76241 Šiauliai, Lithuania

E-mail: aidas.balciunas@mif.vu.lt; a.rimkeviciene@svako.lt

In [3], Mishou obtained a joint universality theorem for the Riemann and Hurwitz zeta-functions. A discrete version of Mishou's theorem was given in [2].

Let $\mathbf{a} = \{a_m : m \in \mathbb{N}\}$ and $\mathbf{b} = \{b_m : m \in \mathbb{N}_0\}$ be two periodic sequences of the complex numbers. Moreover, for a fixed $\theta > \frac{1}{2}$, $v_u(m) = \exp\left\{-\left(\frac{m}{u}\right)^\theta\right\}$ and $v_u(m, \alpha) = \exp\left\{-\left(\frac{m+\alpha}{u}\right)^\theta\right\}$, where $0 < \alpha \leq 1$ and $u > 0$. Define two series $\zeta_u(s; \mathbf{a}) = \sum_{m=1}^{\infty} \frac{a_m v_u(m)}{m^s}$ and $\zeta_u(s, \alpha; \mathbf{b}) = \sum_{m=0}^{\infty} \frac{b_m v_u(m, \alpha)}{(m+\alpha)^s}$. Then the later series are absolutely convergent in any half-plane $\sigma > \sigma_0$. These series are certain convolutions with the periodic and periodic Hurwitz zeta-functions, respectively. Let, for positive h_1 and h_2 , $L(\mathbb{P}; \alpha, h_1, h_2, \pi) = \{(h_1 \log p : p \in \mathbb{P}), (h_2 \log p : p \in (m + \alpha)), 2\pi\}$, where \mathbb{P} is the set of all prime numbers. In the report, we will discuss approximation of a pair of analytic functions by shifts of the functions $\zeta_{u_N}(s; \mathbf{a})$ and $\zeta_{u_N}(s, \alpha; \mathbf{b})$ with $u_N \rightarrow \infty$ as $N \rightarrow \infty$. Let \mathcal{K} be the class of compact sets of the strip $\{s \in \mathbb{C} : \frac{1}{2} < \sigma < 1\}$ with connected complements, and $H_0(K)$ and $H(K)$ with $K \in \mathcal{K}$ the classes of continuous nonvanishing and continuous functions on K , respectively, that are analytic in the interior of K . Then the following statement is valid [1].

THEOREM 1. *Suppose that the sequence \mathbf{a} is multiplicative, the set $L(\mathbb{P}; \alpha, h_1, h_2, \pi)$ is linearly independent over the field of rational numbers, and $u_N \rightarrow \infty$ and $u_N \ll N^2$ as $N \rightarrow \infty$. Let $K_1, K_2 \in \mathcal{K}$ and $f_1(s) \in H_0(K_1)$, $f_2(s) \in H(K_2)$. Then, for every $\epsilon > 0$, the limit*

$$\lim_{N \rightarrow \infty} \frac{1}{N+1} \#\left\{0 \leq k \leq N : \sup_{s \in K_1} |\zeta_{u_N}(s + ikh_1; \mathbf{a}) - f_1(s)| < \epsilon_1, \right. \\ \left. \sup_{s \in K_2} |\zeta_{u_N}(s + ikh_2, \alpha; \mathbf{b}) - f_2(s)| < \epsilon_2 \right\}$$

exists and is positive for all but at most countably many $\epsilon_1 > 0$ and $\epsilon_2 > 0$.

REFERENCES

- [1] A. Balčiūnas, M. Jاسas and A. Rimkevičienė. A discrete version of the Mishou theorem related to periodic zeta-Function. *Mathematical Modelling and analysis*, (to appear 2024).
- [2] E. Buivydas and A. Laurinčikas. A discrete version of the Mishou theorem. *Ramanujan J.*, **38** (2):331–347, 2015.
- [3] H. Mishou. The joint value-distribution of the Riemann zeta function and Hurwitz zeta functions. *Lith. Math. J.*, **47** (1):32-47, 2007.

A GLOBAL APPROXIMATION METHOD FOR SECOND-KIND NONLINEAR INTEGRAL EQUATIONS

LUISA FERMO, ANNA LUCIA LAGUARDIA, CONCETTA LAURITA and MARIA GRAZIA
RUSSO

Department of Mathematics, University of Cagliari

Via Ospedale 72, 09124 Cagliari, Italy

E-mail: luisa.fermo@unica.it

Department of Mathematics; computer Science and Economics, University of Basilicata

via dell'Ateneo Lucano 10, Potenza, Italy

E-mail: annalucia.laguardia@unibas.it

E-mail: concetta.laurita@unibas.it

E-mail: mariagrazia.russo@unibas.it

A global approximation method of Nyström type is explored for the numerical solution of a class of nonlinear integral equations of the second kind of the type:

$$f(y) - \int_{-1}^1 k_1(x, y)f(x)dx - \int_{-1}^1 k_2(x, y)h(x, f(x))dx = g(y), \quad y \in [-1, 1],$$

Integral equations of this type have wide applications in problems involving nonlinearities such as heat radiation, heat transfer, acoustic, elasticity, and electromagnetic problems; see [1; 2].

Some of these models are mathematically formulated in terms of boundary value problems having nonlinear boundary conditions, which can be reformulated as the equation under consideration [3; 4].

In applicative contexts, the kernels of the equation are smooth and/or weakly singular. Therefore cases of smooth and weakly singular kernels are both considered. Firstly the solvability of the equation is studied in the space of continuous functions and the smoothness of the solution is stated according to the smoothness of the known functions. In the case of regular kernels, the proposed numerical method uses a Gauss-Legendre rule, whereas in the second case of weakly singular kernels, the method resorts to a product rule based on Legendre nodes. Stability and convergence are proved in subspaces of Sobolev and Zygmund type, equipped with the uniform norm.

REFERENCES

- [1] R. Bialecki, A. J. Nowak. Boundary value problems in heat conduction with nonlinear material and nonlinear boundary conditions. *Appl. Math. Model.*, **5** (6):417–421, 1981.
- [2] M. A. Kelmanson. Solution of nonlinear elliptic equations with boundary singularities by an integral equation method. *J. Comput. Phys.*, **56** (2):224–258, 1984.
- [3] K. Ruotsalainen, W. Wendland. On the boundary element method for some nonlinear boundary value problems. *Numer. Math.*, **53** (:299–314, 1988.
- [4] K. E. Atkinson, G. Chandler. Boundary integral equation methods for solving Laplace's equation with nonlinear boundary conditions: the smooth boundary case. *Math. Comp.*, **55** (192):451–472, 1990.

ON SENSITIVE DEPENDENCE OF SOLUTIONS ON THE INITIAL DATA IN ORDINARY DIFFERENTIAL EQUATIONS

FELIX SADYRBAEV^{1,2}

Institute of Mathematics and Computer Science, University of Latvia

Rainis boulevard 29, LV-1459 Rīga, Latvia

Daugavpils University

13 Vienības street, Daugavpils, LV-5401, Latvia

E-mail: `felix@latnet.lv`

A sensitive dependence of solutions on the initial data is an indication of chaotic behavior. This phenomenon was extensively studied for three-dimensional systems. It appears and plays an important role in many processes of real life. Examples in biology, chemistry, financial mathematics are well known.

In our talk several less-known examples are considered, starting from Duffing-type equations, and including formally very simple ordinary differential equations of order three and four.

Examples are provided, and illustrative materials are supplied.

REFERENCES

- [1] F. Sadyrbaev. On Solutions of the Third-Order Ordinary Differential Equations of Emden-Fowler Type. *Dynamics*, **3** (3):550–562, 2023.
- [2] J.C. Sprott. *Elegant Chaos: Algebraically Simple Chaotic Flows*. World Scientific, 2010.

ON THE MEAN SQUARE OF THE HURWITZ ZETA-FUNCTION IN SHORT INTERVALS

DARIUS ŠIAUČIŪNAS

Vilnius University

Vytauto street 84, LT-76352 Šiauliai, Lithuania

E-mail: `darius.siauciunas@sa.vu.lt`

Let $s = \sigma + it$ be a complex variable, and $0 < \alpha \leq 1$ a fixed parameter. The Hurwitz zeta-function $\zeta(s, \alpha)$ is given, for $\sigma > 1$, by Dirichlet series

$$\zeta(s, \alpha) = \sum_{m=0}^{\infty} \frac{1}{(m + \alpha)^s},$$

and has analytic continuation to the whole complex plane, except for a simple pole $s = 1$.

In the theory of zeta-functions, mean square estimates play an important role. For example, they are the main ingredient in proofs of universality theorems on approximation of analytic functions. A problem of effectivization of universality theorems requires mean square estimates in short intervals.

Our report is devoted to a mean square estimate of the function $\zeta(s, \alpha)$ in short intervals. We will announce the following theorem

THEOREM 1. *Suppose that $\alpha \neq 1, 1/2$, and $1/2 < \sigma < 7/12$ is fixed. Let $T^{27/82} \leq H \leq T^\sigma$. Then, uniformly in H , the estimate*

$$\int_{T-H}^{T+H} |\zeta(\sigma + it, \alpha)|^2 dt \ll_{\sigma, \alpha} H$$

is valid.

Theorem 1 extends a known results for the Riemann zeta-function [1], i.e., for $\zeta(s, 1)$. Note that the case of $\zeta(s, \alpha)$ is different from that of $\zeta(s, 1)$ because $\zeta(s, \alpha)$ has no Euler's product over primes.

REFERENCES

- [1] A. Ivič. *The Riemann Zeta-Function: The Theory of the Riemann Zeta-Function with Applications*. John Wiley & Sons, New York, 1985.

ON ZETA-FUNCTION OF CERTAIN CUSP FORMS

ANTANAS LAURINČIKAS¹ and DARIUS ŠIAUČIŪNAS²

¹*Faculty of Mathematics and informatics, Vilnius University*

Naugarduko street 24, LT-03225 Vilnius, Lithuania

²*Šiauliai Academy, Vilnius University*

Vytauto street 84, LT-76352 Šiauliai, Lithuania

E-mail: antanas.laurincikas@mif.vu.lt, darius.siauciunas@sa.vu.lt

Let $\mathcal{F}(z)$ be a cusp form of weight κ for the full modular group with the Fourier series expansion $\mathcal{F}(z) = \sum_{m=-\infty}^{\infty} c(m)e^{2\pi imz}$. The corresponding zeta-function $\zeta(s, \mathcal{F})$, $s = \sigma + it$, is defined for $\sigma > (\kappa + 1)/2$, by the series

$$\zeta(s, \mathcal{F}) = \sum_{m=1}^{\infty} \frac{c(m)}{m^s},$$

and has analytic continuation to an entire function. We suppose additionally that $\mathcal{F}(z)$ is a normalized Hecke-eigen cusp form. Thus, $\zeta(s, \mathcal{F})$ has the Euler product over prime numbers.

Our report is devoted to universality of the function $\zeta(s, \mathcal{F})$. Let $D_{\mathcal{F}} = \{s \in \mathbb{C} : \kappa/2 < \sigma < (\kappa + 1)/2\}$. Then it is known [1] that the shifts $\zeta(s + i\tau, \mathcal{F})$, $\tau \in \mathbb{R}$, approximate any non-vanishing analytic function defined on $D_{\mathcal{F}}$. In the report, we discuss the density of approximating shifts in short intervals. For this, we use the following conjecture: there exists $\sigma_0 \in (\kappa/2, (\kappa + 1)/2)$ and $0 < \delta < 1$ such that, for $\sigma \in (\kappa/2, \sigma_0]$ and $T^\delta \leq H \leq T$, uniformly in H the estimate

$$\int_{T-H}^{T+H} |\zeta(s + i\tau; \mathcal{F})|^2 dt \ll_{\sigma} H$$

is valid.

Let $\mathcal{K}_{\mathcal{F}}$ be the class of compact sets of $D_{\mathcal{F}}$ with connected complements, and $H_{0\mathcal{F}}(K)$, $K \in \mathcal{K}$, the class of continuous non-vanishing functions on K that are analytic in the interior of K .

THEOREM 1. *Suppose that all above conditions for the form $\mathcal{F}(z)$ are valid. Let $K \in \mathcal{K}_{\mathcal{F}}$, $f(s) \in H_{0\mathcal{F}}(K)$, and $T^\delta \leq H \leq T$. Then, for any $\varepsilon > 0$,*

$$\liminf_{T \rightarrow \infty} \frac{1}{H} \text{meas} \left\{ \tau \in [T, T + H] : \sup_{s \in K} |f(s) - \zeta(s + i\tau; \mathcal{F})| < \varepsilon \right\} > 0.$$

Moreover, “lim inf” can be replaced by “lim” for all but at most countably many $\varepsilon > 0$.

Proof of Theorem 1 will appear in [2].

REFERENCES

- [1] A. Laurinčikas and K. Matsumoto. The universality of zeta-functions attached to certain cusp forms. *Acta Arith.*, **98** (4):345–359, 2001.
- [2] A. Laurinčikas and D. Šiaučiūnas. On universality in short intervals for zeta-functions of certain cusp forms. *Math. Slovaca*, (accepted).

MATHEMATICS OF EXTREMES IN COASTAL AND MARINE SCIENCE

TARMO SOOMERE

Tallinn University of Technology

Ehitajate tee 5, 19086 Tallinn, Estonia

E-mail: tarmo.soomere@taltech.ee

Marine-driven hazards are often created by a variety of components that have not only different nature, temporal and spatial scales, but also exhibit radically different statistical properties of their extremes. For example, extreme water levels are jointly created by (i) storm surges that follow an exponential distribution, (ii) the volume of excess water in the Baltic Sea that follows a normal distribution, and (iii) wave-generated local water level setup that often obeys an inverse Gaussian (Wald) distribution.

The most dangerous water levels follow a generalized extreme value (GEV) or (generalized) Pareto distribution. The properties of both extreme water levels and these distributions vary substantially in different coastal segments of the eastern Baltic Sea, providing not only delicate information about a variety of driving mechanisms of waves and water level but also unique opportunities for insight into future of marine and coastal extremes.

I intend to narrate a story about how extreme water levels are born in the Baltic Sea [1], how it is possible to separate their drivers and identify which distributions they follow [2], which important message is provided via the simplest (trend) analysis of increasing water level maxima [3] and how these trends coexist with the contradicting narratives of (non)stationarity, changing climate [4] and human impacts [5]. Further on I delve deeper into the present and future of extremely high and low water levels on the shores of Estonia, Latvia and Lithuania [6]. Finally, I intend to provide a small selection of unexpected features of extremes of various kinds in our coastal regions [7].

REFERENCES

- [1] T. Soomere and K. Pindsoo. Spatial variability in the trends in extreme storm surges and weekly-scale high water levels in the eastern Baltic Sea. *Continental Shelf Research*, **115**, 53–64, 2016. doi: 10.1016/j.csr.2015.12.016.
- [2] T. Soomere, M. Eelsalu, A. Kurkin and A. Rybin. Separation of the Baltic Sea water level into daily and multi-weekly components. *Continental Shelf Research*, **103**, 23–32, 2014. doi: 10.1016/j.csr.2015.04.018.
- [3] K. Pindsoo and T. Soomere. Basin-wide variations in trends in water level maxima in the Baltic Sea. *Continental Shelf Research*, **193**, 104029, 2020. doi: 10.1016/j.csr.2019.104029.
- [4] N. Kudryavtseva, T. Soomere and R. Männikus. Non-stationary analysis of water level extremes in Latvian waters, Baltic Sea, during 1961–2018. *Natural Hazards and Earth Systems Sciences*, **21**, 1279–1296, 2021. doi: 10.5194/nhess-21-1279-2021.
- [5] M. Reckermann, A. Omstedt, T. Soomere, J. Aigars, N. Akhtar, M. Beldowska, J. Beldowski, T. Cronin, M. Czub, M. Eero, K.P. Hyytiäinen, J.-P. Jalkanen, A. Kiessling, E. Kjellström, K. Kuliński, X. Guo Larsen, M. McCrackin, H.E.M. Meier, S. Oberbeckmann, K. Parnell, C. Pons-Seres de Brauwer, A. Poska, J. Saarinen, B. Szymczycha, E. Undeman, A. Wörman and E. Zorita. Human impacts and their interactions in the Baltic Sea region. *Earth Systems Dynamics*, **13**, 1–80, 2022. doi: 10.5194/esd-13-1-2022.
- [6] K. Viigand, M. Eelsalu and T. Soomere. Quantifying exposedness of the eastern Baltic Sea shores with respect to extremely high and low water levels. *Coastal Engineering*, under review.
- [7] T. Soomere. Climate change and coastal processes in the Baltic Sea. In: *Oxford Encyclopedia of Climate Science*, online publication 2024, doi: 10.1093/acrefore/9780190228620.013.897.

COLLOCATION BASED APPROXIMATIONS FOR FRACTIONAL BOUNDARY VALUE PROBLEMS

HANNA BRITT SOOTS

Institute of Mathematics and Statistics, University of Tartu, Estonia

Narva mnt 18, Tartu, Tartumaa, Estonia

E-mail: hanna.britt.soots@ut.ee

We construct a high order method for the numerical solution of fractional weakly singular integro-differential equations in the form

$$(D_{\text{Cap}}^\alpha y)(t) + h(t)y(t) + \int_0^t L_\kappa(t, s)y(s)ds = f(t), \quad 0 \leq t \leq b, \quad 0 < b < \infty, \quad (1)$$

subject to the conditions

$$a_{11}y(0) + a_{12}y(b_1) = \gamma_1; \quad a_{21}y'(0) + a_{22}y(b_1) = \gamma_2. \quad (2)$$

The problem is reformulated as an integral equation of the second kind with respect to $z = D_{\text{Cap}}^\alpha y$, the Caputo fractional derivative of y of order α , with $1 < \alpha < 2$, where y is the solution of the original problem. Using this reformulation, the regularity properties of both y and its Caputo derivative z are studied. Based on this information a piecewise polynomial collocation method is developed for finding an approximate solution z_N of the reformulated problem. Using z_N , an approximation y_N for y is constructed and a detailed convergence analysis of the proposed method is given. In particular, the attainable order of convergence of the proposed method for appropriate values of grid and collocation parameters is established. To illustrate the performance of our approach, results of some numerical experiments are presented.

REFERENCES

- [1] H.B. Soots, K. Lätt, A. Pedas. Collocation based approximations for a class of fractional boundary value problems. *Mathematical Modelling and Analysis*, **28** (2).

DISCRETE STURM–LIOUVILLE PROBLEM FOR TWO-DIMENSIONAL ELLIPTIC EQUATION WITH THE MULTIPLE INTEGRAL IN NONLOCAL CONDITION

ARTŪRAS ŠTIKONAS¹ and MIFODIJUS SAPAGOVAS²

¹*Institute of Applied Mathematics, Vilnius University*

Naugarduko str. 24, LT-03225 Vilnius, Lithuania

²*Institute of Data Science and Digital Technologies, Vilnius University*

Akademijos str. 4, Vilnius LT-08412, Lithuania

E-mail: arturas.stikonas@mif.vu.lt; mifodijus.sapagovas@mif.vu.lt

We consider the following Sturm-Liouville Problem (SLP) with nonlocal boundary condition (NBC) involving multiple integral [2]:

$$L(u) := -\frac{\partial^2 u}{\partial x^2} - \frac{\partial^2 u}{\partial y^2} = \lambda u, \quad (x, y) \in \Omega \subset \mathbb{R}^2, \quad (1)$$

$$u(x, y) = \int_{\Omega} k(x, y, \xi, \eta) u(\xi, \eta) d\xi d\eta, \quad (x, y) \in \partial\Omega, \quad (2)$$

where $\Omega := \Omega_x \times \Omega_y$, $\Omega_x := \{x: 0 < x < a\}$, $\Omega_y := \{y: 0 < y < b\}$. We will assume that the function k satisfies the condition $\int_{\Omega} |k(x, y, \xi, \eta)| d\xi d\eta \leq \varrho < 1$. The finite difference method for the linear two-dimensional parabolic equation with the NBC (2) in the square domain $\Omega = [0, 1]^2$ was studied in [1].

Let us write the Finite-Difference Scheme for SLP (1)–(2):

$$L^h(U) := -\delta_x^2 U - \delta_y^2 U = \lambda U, \quad (x_i, y_j) \in \omega^h, \\ U_{ij} = [K_{ij}, U]^{\text{tr}}, \quad (x_i, y_j) \in \partial\bar{\omega}^h,$$

and we use two-dimensional trapezoidal rule $[K_{ij}, U]^{\text{tr}}$ where $K_{ij}^{kl} = k(x_i, y_j, x_k, y_l)$, $(x_i, y_j) \in \partial\bar{\omega}^h$, $(x_k, y_l) \in \bar{\omega}^h$ for approximation of integral in BC. We suppose that $\max_{x_{ij} \in \partial\bar{\omega}^h} [|K_{ij}|, 1]^{\text{tr}} \leq \rho < 1$.

The main aims of our study are the investigation SLP for various cases of kernel K . The main difficulty of this problem is that in the non-classical case we cannot use the method of separation of variables and decompose the problem into one-dimensional problems. Numerical experiments show that the spectrum of this problem is similar to the spectrum of corresponding one-dimensional problems. For example, all eigenvalues are real.

REFERENCES

- [1] Y. Lin, S. Xu and H.-M. Yin. Finite difference approximation for a class of non-local parabolic equations. *Internat. J. Math. & Math. Sci.*, **1997** 20(1):147–163, 1997. <https://doi.org/10.1155/S0161171297000215>
- [2] Y. Wang. Solutions to nonlinear elliptic equations with a nonlocal boundary condition. *Electron. J. Differential Equations*, **2002** (05):1–16, 2002. <https://ejde.math.txstate.edu/Volumes/2002/05/abstr.html>

FINITE DIFFERENCE METHOD FOR ELLIPTIC EQUATION WITH TWO MULTIPLE INTEGRAL CONDITIONS

OLGA ŠTIKONIENĖ¹ and MIFODIJUS SAPAGOVAS²

¹*Institute of Applied Mathematics, Vilnius University*

Naugarduko 24, LT-03225, Vilnius, Lithuania

²*Institute of Data Science and Digital Technologies, Vilnius University*

Akademijos 4, LT-08663, Vilnius, Lithuania

E-mail: olga.stikoniene@mif.vu.lt; mifodijus.sapagovas@mif.vu.lt

We consider a finite difference approximation to the solution of the nonlocal boundary value problem for two-dimensional Poisson equation

$$-\frac{\partial^2 u}{\partial x^2} - \frac{\partial^2 u}{\partial y^2} = f(x, y), \quad (x, y) \in \Omega \subset \mathbb{R}^2$$

with nonlocal conditions including multiple integrals

$$\begin{aligned} u(x, 0) &= \gamma_0 \int_{\Omega} u(\xi, \eta) d\xi d\eta + g_0(x), \quad x \in \Omega_x, \\ u(x, L_y) &= \gamma_1 \int_{\Omega} u(\xi, \eta) d\xi d\eta + g_1(x), \quad x \in \Omega_x, \\ u(0, y) &= g_2(y), \quad u(L_x, y) = g_3(y), \quad y \in \Omega_y, \end{aligned}$$

where $\Omega := \Omega_x \times \Omega_y$, $\Omega_x := \{x: 0 < x < L_x\}$, $\Omega_y := \{y: 0 < y < L_y\}$.

We evaluate double integrals using Trapezoidal rule in two dimensions. The Laplace operator is approximated in a standard way on a five-point stencil.

The main aim is to find the conditions for the parameters γ_0 and γ_1 under which unique solution exists.

ON SIMULATION WITH PARTICLE MOTION TRACKING FOR BIOMASS EXTRACTION PROCESS

ULDIS STRAUTIŅŠ^{1,2}, MAKSIMS MARINAKI^{1,2}

Institute of Mathematics and Computer Science¹, University of Latvia²

Raiņa bulvāris 29, Rīga, LV-1459, Latvija¹

Jelgavas iela 3, Rīga, LV-1004, Latvija²

E-mail: `uldis.strautins@lu.lv`

E-mail: `maksims.marinaki@lu.lv`

The extraction process of active components from biomass is based on filtration of the solvent through a filter cake of biomass. We focus on the simulation of the evolution of the filter cake. The computational domain is separated into two parts: the fluid part $\Omega_F(t)$ and the filter cake part $\Omega_C(t)$ with a time dependent interface separating the two. The filter cake can grow by adhesion of new biomass particles carried by the flow in Ω_F .

As a result, one obtains a coupled system of Navier-Stokes and Darcy equations

$$u_t + u \cdot \nabla u = \nu \Delta u - \nabla p + f, \quad \Omega_F;$$

$$u = -K \nabla p + f, \quad \Omega_C;$$

$$\nabla \cdot u = 0, \quad \Omega_F \cup \Omega_C.$$

On the interface Beavers-Joseph conditions are imposed [1].

The particles are assumed to be round and influenced by hydrodynamic drag force and gravity. The particles are assumed to stick to the filter cake upon collision. As a result, the filter cake domain grows in time.

The equations are implemented using a finite volume scheme on a voxel mesh extending a previous solver [2]. The interface between the Darcy and Navier-Stokes domains is not resolved fully and changes in discrete steps corresponding to the voxel size.

REFERENCES

- [1] M. Discacciati, A. Quarteroni. Navier-Stokes/Darcy coupling: Modeling, analysis and numerical approximation. *Rev. Mat. Complut.*, **22** (2):315–426, 2009.
- [2] U. Strautins, M. Marinaki. On modelling of three dimensional flow in extraction of biologically active substances from plants. *Eng. Rural Dev.*, **22** 204 – 209, 2023.

A GENERAL COLLOCATION ANALYSIS FOR WEAKLY SINGULAR VOLTERRA INTEGRAL EQUATIONS WITH VARIABLE EXPONENT

MARTIN STYNES

Beijing Computational Science Research Center

Haidian District, Beijing 1000193, China

E-mail: m.stynes@csrc.ac.cn

Piecewise polynomial collocation of weakly singular Volterra integral equations (VIEs) of the second kind has been extensively studied in the literature, where integral kernels of the form $(t - s)^{-\alpha}$ for some constant $\alpha \in (0, 1)$ are considered. Variable-order fractional-derivative differential equations currently attract much research interest, and in Zheng and Wang *SIAM J. Numer. Anal.* 2020 such a problem is transformed to a weakly singular VIE whose kernel has the above form with variable $\alpha = \alpha(t)$, then solved numerically by piecewise linear collocation, but it is unclear whether this analysis could be extended to more general problems or to polynomials of higher degree. In the present paper the general theory (existence, uniqueness, regularity of solutions) of variable-exponent weakly singular VIEs is developed using novel techniques. These results then underpin an error analysis of collocation methods where piecewise polynomials of any degree can be used. This error analysis is also novel — it makes no use of the usual resolvent representation, which is a key technique in the error analysis of collocation methods for VIEs in the current research literature. Furthermore, all the above analysis for a scalar VIE can be extended to certain nonlinear VIEs and to systems of VIEs. The sharpness of the theoretical error bounds obtained for the collocation methods is demonstrated by numerical examples.

This is joint work with Professor Hui Liang.

REFERENCES

- [1] H. Liang and M. Stynes. Article in journal. *IMA J. Numer. Anal.*,(to appear). doi: <https://doi.org/10.1093/imanum/drad072>

ON DERIVATIVE SAMPLING AND CONVERGENCE IN VARIATION OF GENERALIZED SHANNON SAMPLING SERIES

GERT TAMBERG and MARIA ZELTSER

Tallinn University of Technology

Ehitajate tee 5, 19086 Tallinn, Estonia

E-mail: gert.tamberg@taltech.ee

Tallinn University

Narva mnt 25, 10120 Tallinn, Estonia

E-mail: maria.zeltser@tlu.ee

In this presentation we consider generalized operators for approximating derivatives and also in the case of functions with bounded variation.

We take the formula, connecting generalized sampling operators $G_w^{\bar{x}}$ and sampling Kantorovich operators K_w^x , presented in [2] by D. Costarelli and G. Vinti in form

$$(K_w^x f)(x) = (G_w^{\bar{x}} F)' \left(x + \frac{1}{2w} \right) \quad (1)$$

and give it a more simpler form. To do that, we use generalized Kantorovich-type (or Durrmeyer) sampling operators, we introduced in [4] and related kernels, introduced in [3]. This approach allows us to study derivative sampling. We give estimates of order of approximation in terms of moduli of smoothness.

An overview about sampling Kantorovich operators in BV-spaces is given in [1]. We use generalized Kantorovich-type sampling operators in BV-spaces and give estimates of order of approximation in terms of corresponding moduli of smoothness.

REFERENCES

- [1] L. Angeloni, and G. Vinti. Multidimensional sampling-Kantorovich operators in BV-spaces. *Open Math.*, **21** (1):20220573, 2023.
- [2] D. Costarelli, and G. Vinti. Inverse results of approximation and the saturation order for the sampling Kantorovich series. *Journal of Approximation Theory*, **242** (1):64-82, 2019.
- [3] A. Kivinukk and T. Metsmägi. The Variation Detracting Property of Some Shannon Sampling Series and Their Derivatives. *Sampling Theory in Signal and Image Processing*, **13** (2):189-206, 2014.
- [4] O. Orlova, G. Tamberg. On approximation properties of generalized Kantorovich-type sampling operators. *Journal of Approximation Theory*, **201** (1):73-86, 2016.

APPROXIMATION OF ANALYTIC FUNCTIONS BY SHIFTS OF THE PERIODIC ZETA-FUNCTIONS

MONIKA TEKORĖ

Vilnius University

Naugarduko str. 24, LT-03225 Vilnius, Lithuania

E-mail: monika.tekore@mif.stud.vu.lt

Let $s = \sigma + it$ be a complex variable, and $\mathbf{a} = \{a_m : m \in \mathbb{N}\}$ a periodic sequence of complex numbers with minimal period $q \in \mathbb{N}$. The periodic zeta-function $\zeta(s; \mathbf{a})$ is defined, for $\sigma > 1$, by the series

$$\zeta(s; \mathbf{a}) = \sum_{m=1}^{\infty} \frac{a_m}{m^s},$$

and is meromorphically continued to the whole complex plane with a possible simple pole at the point $s = 1$. In the report, we consider approximation of analytic functions by shifts $\zeta(s + i\varphi(t); \mathbf{a})$ with a certain function $\varphi(t)$ and multiplicative sequence \mathbf{a} .

We will discuss the following problems.

1. Approximation of analytic functions by discrete shifts $\zeta(s + iht_k; \mathbf{a})$, $h > 0$, $k \in \mathbb{N}$, where $\{t_k\}$ is a sequence of Gram points [4].

2. Joint approximation of collections of analytic functions by shifts $(\zeta(s + i\gamma_1(\tau); \mathbf{a}_1), \dots, \zeta(s + i\gamma_r(\tau); \mathbf{a}_r))$, where $\gamma_1(\tau), \dots, \gamma_r(\tau)$ are increasing to infinity continuously differentiable functions [1].

3. Joint approximation of collections of analytic functions by shifts $(\zeta(s + ih_1\gamma_k; \mathbf{a}_1), \dots, \zeta(s + ih_r\gamma_k; \mathbf{a}_r))$, $h_j > 0$, where $\{\gamma_k : k \in \mathbb{N}\}$ is a sequence of non-trivial zeros of the Riemann zeta-function [2].

4. Approximation of analytic functions by shifts of compositions $F(\zeta(s + ih_1\gamma_k; \mathbf{a}_1), \dots, \zeta(s + ih_r\gamma_k; \mathbf{a}_r))$, where $F : H^r(D) \rightarrow H(D)$ is a certain continuous operator, and $H(D)$ denotes the space of analytic functions on $D = \{s \in \mathbb{C} : 1/2 < \sigma < 1\}$ [3].

The above results form the doctoral dissertation of the author.

REFERENCES

- [1] A. Laurinčikas and M. Tekorė. Joint universality of periodic zeta-functions with multiplicative coefficients. *Nonlinear Analysis: Modelling and Control*, **25** (5):860–883, 2020.
- [2] A. Laurinčikas, D. Šiaučiūnas and M. Tekorė. Joint universality of periodic zeta-functions with multiplicative coefficients. II. *Nonlinear Analysis: Modelling and Control*, **26** (3):550–563, 2021.
- [3] D. Šiaučiūnas, R. Šimėnas and M. Tekorė. Approximation of analytic functions by shifts of certain compositions. *Mathematics*, **9** (20):2583, 2021.
- [4] D. Šiaučiūnas and M. Tekorė. Gram points in the universality of the Dirichlet series with periodic coefficients. *Mathematics*, **11** (22):4615, 2023.

TIME-FREQUENCY ANALYSIS IN EUCLIDEAN SPACES

VILLE TURUNEN

Aalto University

Department of Mathematics and Systems Analysis, P.O. Box 11100, FI-00076 Aalto, Finland

E-mail: `ville.turunen@aalto.fi`

Time-frequency analysis can be described as Fourier analysis simultaneously both in time and in frequency. Its origins are in quantum mechanics, in signal processing, and in pseudo-differential operators. A time-frequency transform is a sesquilinear mapping from a family of test functions in time to functions in the time-frequency plane. The class of time-frequency transforms is further restricted by imposing conditions stemming from basic transformations of signals and those which an idealized energy density could satisfy. In [1] with Vesa Vuojamo and Heikki Orelma, we characterized time-frequency transforms in Euclidean spaces in terms of the corresponding pseudo-differential operator quantizations and integral kernel conditions. In the talk, I will also show how to generalize time-frequency analysis to those locally compact groups that allow a nice-enough Fourier transform: wide families of topological groups can be treated, including all the compact groups and all the locally compact commutative groups, and more.

REFERENCES

- [1] Vesa Vuojamo, Ville Turunen, Heikki Orelma. Time-frequency analysis in \mathbb{R}^n . *Journal of Fourier Analysis and Applications*, **28** (6): 1–38, 2022.

THE FUČÍK TYPE PROBLEM WITH NONLOCAL TWO-POINT BOUNDARY CONDITIONS

SIGITA URBONIENĒ¹, NATALIJA SERGEJEVA²

¹ *Vytautas Magnus University*

Universiteto str. 10, Akademija, Kaunas region, LT-46265 Lithuania

² *Latvia University of Life Sciences and Technologies*

Liela str. 2, Jelgava, LV-3001 Latvia

E-mail: sigita.urboniene@vdu.lt, natalija.sergejeva@lbtu.lv

Let us investigate the Fučík problem

$$x'' = -\mu x^+ + \lambda x^-, \quad (1)$$

with one classical

$$x(0) = 0, \quad (2)$$

and other nonlocal two-point boundary condition

$$x(1) = \gamma x(\xi), \quad (3)$$

$$x(1) = \gamma x'(\xi), \quad (4)$$

here $\xi = \frac{m}{n} \in (0, 1)$, $\gamma \in \mathbb{R}$, m and n ($0 < m < n$) are positive coprime integer numbers.

Also we analyze the Fučík problem (1) with condition

$$x'(0) = 0 \quad (5)$$

in the left boundary, as well as with nonlocal conditions (3) – (4) in the right boundary. The aim of investigation is to analyze main properties of the nonlocal problem spectrum and compare them with classical Fučík spectrum. The idea of boundary conditions was taken from the work [1], where the Sturm-Liouville equation

$$x'' = -\lambda x$$

was analyzed with boundary conditions (2) - (5).

Some of the results are the logical continuation and generalization of previous authors' investigations [2; 3; 4].

REFERENCES

- [1] S. Pečiulytė, A. Štikonas. On positive eigenfunctions of Sturm–Liouville problem with nonlocal two-point boundary condition. *Math. Model. Anal.*, **12** (2):2015–226, 2007.
- [2] N. Sergejeva. The regions of solvability for some three point problem. *Math. Model. Anal.*, **18** (2):191–203, 2013.
- [3] N. Sergejeva. The Fučík spectrum for some boundary value problem. In: *Proc. of IMCS of University of Latvia*, 14, 65–75. 2014
- [4] N. Sergejeva. On some Fučík type problem with nonlocal boundary condition. In: *Proc. of IMCS of University of Latvia*, 19, 57–64. 2019.

OPERATORS OF FUZZY MATHEMATICAL MORPHOLOGY

KSENIJA VARFOLOMEJEVA¹ and ALEXANDER ŠOSTAK^{1,2}

¹ *Department of Mathematics, University of Latvia*

Jelgavas iela 3, Rīga, LV-1004, Latvia

² *Institute of Mathematics and CS, University of Latvia*

Raiņa bulvāris 29, Rīga, LV-1459, Latvia

E-mail: ksenija.varfolomejeva@lu.lv, aleksandrs.sostaks@lu.lv

Mathematical morphology arose from the needs of practical geology in the 1970s. However soon, fuzzy morphology theory found important applications in other fields, particularly in image processing, and is now widely studied and applied by many researchers. Mathematical morphology theory within fuzzy sets was firstly introduced in the paper by De Baets et al. [1]. As in "classical" approach, the fuzzy mathematical morphology was based on two operators - dilation \mathcal{D} and erosion \mathcal{E} are induced on the linear space by a fuzzy structuring element B .

A lot of work has been done in the study of fuzzy morphological operators on Euclidean space and many applications have been found in the study of different practical problems. Taking as a basis a pair of operators that behave like dilation and erosion lead to various abstract, in particular, algebraic approaches to the subject of mathematical morphology. In turn, this made it possible to develop a categorical view of mathematical morphology, see, e.g. [2], [3], [4] et al. However, the disadvantage of these abstract approaches is that the role of the structural element is practically lost.

As far as we know, there was no much work done to develop "classical"(fuzzy) mathematical morphology theory within the framework of category theory. This in turn essentially restricts the possibility of considering relations between (fuzzy) morphological spaces, their transformations, products and direct sums of (fuzzy) morphological spaces, etc.

The purpose of our talk is to present a category containing "classical" (fuzzy) morphological spaces realized in the spirit of the article [1]. This provides us the flexibility to deal with transformations of fuzzy morphological spaces, which in turn can enrich the use in practical applications. As the basis for this category, we take structured additive groups $(X, S, +_X, 0_X)$ and $(Y, T, +_Y, 0_Y)$ as objects and certain L -fuzzy relations $R : X \times Y \rightarrow L$ between them as morphisms.

REFERENCES

- [1] B. De Baets, E.E. Kerre, M. Gupta. The fundamentals of fuzzy mathematical morphology Part I: basic concepts. *International J. of General Systems*, **23** 155–171, 1995.
- [2] H. Heijmans, C. Ronse. The algebraic basis for mathematical morphology: dilation and erosion. *Computer Vision, Graphics and Image Processing*, **50** 245–295, 1990.
- [3] A. Šostak, I. Uljane. On two categories of many-level morphological spaces. *Studies in Computational Intelligence*, **955** 207–217, 2022.
- [4] N. Madrid, M. Ojeda-Aciego, J. Medina, I. Perfilieva. L-fuzzy relational mathematical morphology based on adjoint triples. *Information Sciences*, **474** 75–89, 2019.

APPROXIMATE SOLUTIONS TO LINEAR FRACTIONAL INTEGRO-DIFFERENTIAL EQUATIONS

MIKK VIKERPUUR

Institute of Mathematics and Statistics, University of Tartu

Narva 18, Tartu, 51009, Estonia

E-mail: mikk.vikerpuur@ut.ee

We consider a class of fractional weakly singular integro-differential equations

$$\begin{aligned} (D_{Cap}^{\alpha_2} y)(t) + d_1(t) (D_{Cap}^{\alpha_1} y)(t) + d_0(t)y(t) + \int_0^t (t-s)^{-\kappa_0} K_0(t,s)y(s)ds \\ + \int_0^t (t-s)^{-\kappa_1} K_1(t,s)(D_{Cap}^{\theta} y)(s)ds = f(t), \quad 0 \leq t \leq b, \end{aligned} \quad (1)$$

subject to boundary conditions

$$a_i y^{(i)}(0) + b_i y^{(i)}(b) = \gamma_i, \quad a_i, b_i, \gamma_i \in \mathbb{R}, \quad i = 0, \dots, n-1. \quad (2)$$

Here D_{Cap}^{δ} is the Caputo differential operator of order $\delta > 0$ and $n := \lceil \alpha_2 \rceil$ is the smallest integer greater or equal to the fractional order α_2 . We assume that

$$0 < \alpha_1 < \alpha_2 \leq n, \quad \theta \in (0, \alpha_2), \quad \kappa_0, \kappa_1 \in [0, 1)$$

and that the given functions d_0, d_1, K_0, K_1 and f are continuous on their respective domains.

On the basis of [1] we study the existence, uniqueness and regularity of the solution y to problem (1)–(2) and show that under suitable conditions this problem can be reformulated as a Volterra integral equation of the second kind with respect to the fractional derivative $D_{Cap}^{\alpha_2} y$. We regularize the solution of this integral equation with the use of a suitable smoothing transformation and construct a numerical solution to the transformed integral equation by applying a piecewise polynomial collocation method on a mildly graded or uniform grid. We show the convergence of the proposed algorithm and present global superconvergence results for a class of specific collocation parameters. Finally, we complement the theoretical results with some numerical examples.

This is a joint work with Arvet Pedas.

REFERENCES

- [1] A. Pedas and M. Vikerpuur. Spline Collocation for Multi-Term Fractional Integro-Differential Equations with Weakly Singular Kernels. *Fractal and Fractional*, **5(3)** 90, 2021.

ON A SOLVABILITY OF A THREE-PARAMETER PROBLEM ARISING IN HEAT TRANSFER

ARMANDS GRITSANS¹ and INARA YERMACHENKO²

Daugavpils University, Institute of Life Sciences and Technology

Parādes str. 1^a, Daugavpils, LV-5400, Latvia

E-mail: ¹armands.gricans@du.lv, ²inara.jermachenko@du.lv

A convective flow in a tall vertical annulus of radii R and 1, in a case when the walls of the annulus are maintained at the same temperature, can be described by the nonlinear boundary value problem

$$T'' + \frac{1-\alpha}{r}T' + F e^T = 0, \quad T(R) = 0 = T(1), \quad (1)$$

where α is a real number, F is a positive number and $R \in (0, 1)$. We study the existence of the positive solutions $T(r)$ of problem (1) depending on the values of the parameters α , F and R .

By using the technique described in [1] and based on the Krasnosels'kiĭ-Guo fixed point theorem of cone expansion and compression, we prove the following theorem.

THEOREM 1. *Suppose that α and R are real numbers and $R \in (0, 1)$. There exists a positive number F^* such that for every $F \in (0; F^*)$ the problem (1) has two positive solutions \underline{T} and \overline{T} with $\|\underline{T}\| < 1 < \|\overline{T}\|$.*

The results obtained allow us to formulate a number of statements about solvability of the problem (1) depending on the values of one of the parameters α , F or R , if other two parameters are fixed.

The results of present study can be used for design of bioreactor systems.

REFERENCES

- [1] A. Gritsans, A. Kolyshkin, F. Sadyrbaev, and I. Yermachenko. On the stability of a convective flow with nonlinear heat sources. *Mathematics*, **2023** 11(18): 3895.
- [2] A. Gritsans, A. Kolyshkin, D. Ogorelova, F. Sadyrbaev, I. Samuilik, and I. Yermachenko. Solutions of nonlinear boundary value problem with applications to biomass thermal conversion. In: *Proc. of 20th Intern. Scientific Conference "Engineering for rural development", Jelgava, Latvia, 2021*, 837-842.
- [3] D. Guo and V. Lakshmikantham. *Nonlinear problems in abstract cones*. Academic Press, 1988.
- [4] G. Infante. *A short course on positive solutions of systems of ODEs via fixed point index*. <https://arxiv.org/abs/1306.4875>, 2017.

Index

- Šņeps-Šņeppe, M., 48
Šostak, A., 70
Štikonas, A., 13, 62
Čiegis, R., 17
Šiaučiūnas, D., 25, 58, 59
Štikonienė, O., 63
- Arriagada, W., 9
Atslega, S., 10
Avitabile, D., 46
- Bajārs, J., 11, 35
Balčiūnas, A., 12, 55
Bingelė, K., 13
Braš, M., 14
Brugnano, L., 15
- Chemetov, N., 46
Christian, S. R., 16
Coelho, C., 22
Costa, M. F. P., 22
- Dapkūnas, B., 18
Diethelm, K., 19
Dzenis, M. G., 20
- Fatone, L., 24
Fermo, L., 21, 56
Ferrás, L. L., 22
Ford, N. J., 23
- Garbaliuskienė, V., 25
Gerges, H., 26, 27
Gritsans, A., 72
Guariglia, E., 28
- Hämarik, U., 36
- Jackiewicz, Z., 29
Janikashvili, N., 34
Janno, J., 26
Jasas, M., 30
Junghanns, P., 44
Juodagalvytė, R., 31
- Kaklauskas, L., 33
Kalichava, Z., 34
Kalvāns, D., 35
Kangro, U., 36
Kaulakytė, K., 37
Kačinskaitė, R., 32
Kirsiaed, E., 38
Kopteva, N., 39
- Kozera, R., 40
Kozulinas, N., 41
- Lätt, K., 42
Laguardia, A. L., 56
Laurinčikas, A., 25, 43, 59
Laurita, C., 44, 56
Lillemäe, M., 45
Lima, P., 46
- Macaitienė, R., 27, 47
Marinaki, M., 48, 64
Mezzanotte, D., 21
Mikalauskaitė, T., 49
- Occorsio, D., 21
Odisharia, K., 34
Odisharia, V., 34
Oja, P., 38
- Prokopenya, A., 50
Pupalaigė, K., 51
- Ragulskis, M., 52
Raus, T., 53
Rebelo, M., 54
Reinfelds, A., 16
Rimkevičienė, A., 55
Russo, M. G., 56
- Sadyrbaev, F., 10, 57
Sapagovas, M., 51, 62, 63
Sergejeva, N., 69
Simina, A., 11
Soomere, T., 60
Soots, H.-B., 61
Strautiņš, U., 64
Stynes, M., 65
- Tamberg, G., 66
Tekorė, M., 67
Turunen, V., 68
- Urbonienė, S., 69
- Varfolomejeva, K., 70
Vikerpuur, M., 71
- WilkoŁazka, M., 40
- Yermachenko, I., 72
- Zeltser, M., 66